

# Generalisation of splitting methods based on modified potentials to nonlinear evolution equations of parabolic and Schrödinger type

S. Blanes, F. Casas, C. González, M. Thalhammer<sup>1</sup>

---

## Abstract

The present work is concerned with the extension of modified potential operator splitting methods to nonlinear evolution equations of Schrödinger and parabolic type. Moreover, an invariance principle is deduced that has a significant impact on the efficient realisation of the resulting modified operator splitting methods for nonlinear Schrödinger equations.

Numerical illustrations for the time-dependent Gross–Pitaevskii equation in the physically most relevant case of three space dimensions and for its parabolic counterpart related to ground state and excited state computations confirm the benefits of the proposed fourth-order modified operator splitting method in comparison with standard splitting methods.

The presented results are novel and of particular interest from both, a theoretical perspective to inspire future investigations of modified operator splitting methods for other classes of nonlinear evolution equations and a practical perspective to advance the reliable and efficient simulation of Gross–Pitaevskii systems in real and imaginary time.

*Key words:* Nonlinear evolution equations, Parabolic problems, Schrödinger equations, Gross–Pitaevskii systems, Geometric time integration, Operator splitting methods, Fourier spectral method, Stability, Convergence, Efficiency.

---

<sup>1</sup> Addresses: Sergio Blanes, Universitat Politècnica de València, Instituto de Matemática Multidisciplinar, 46022 Valencia, Spain. Fernando Casas, Universitat Jaume I, IMAC and Departament de Matemàtiques, 12071 Castellón, Spain. Cesáreo González, Universidad de Valladolid, Departamento de Matemática Aplicada, 47011 Valladolid, Spain. Mechthild Thalhammer, Leopold–Franzens–Universität Innsbruck, Institut für Mathematik, 6020 Innsbruck, Austria. Websites: [personales.upv.es/serblaza](http://personales.upv.es/serblaza), [www.gicas.uji.es/fernando.html](http://www.gicas.uji.es/fernando.html), [www.imuva.uva.es/en/investigadores/33](http://www.imuva.uva.es/en/investigadores/33), [techmath.uibk.ac.at/mecht](http://techmath.uibk.ac.at/mecht). Email addresses: [serblaza@imm.upv.es](mailto:serblaza@imm.upv.es), [fernando.casas@uji.es](mailto:fernando.casas@uji.es), [ome@am.uva.es](mailto:ome@am.uva.es), [mechthild.thalhammer@uibk.ac.at](mailto:mechthild.thalhammer@uibk.ac.at).

## 1 Introduction

**Scope of applications.** A wide range of relevant applications in sciences includes the numerical integration of initial value problems for nonlinear evolution equations. In many cases, the function defining the right-hand side comprises two or more parts

$$\begin{cases} \frac{d}{dt} u(t) = F_1(u(t)) + F_2(u(t)), \\ u(0) = u_0, \quad t \in [0, T]. \end{cases} \quad (1)$$

As prominent instances, we highlight time-dependent nonlinear Schrödinger equations, more specifically, Gross–Pitaevskii equations that arise in the description of Bose–Einstein condensation, see [15,23]. For a comprehensive overview of the underlying principles of quantum theory, we refer to [22].

**Nonlinear Schrödinger equation (Gross–Pitaevskii equation).** A fundamental model for the nonlinear dynamics of a single Bose–Einstein condensate reads as

$$\begin{cases} i \partial_t \Psi(x, t) = -\Delta \Psi(x, t) + V(x) \Psi(x, t) + \vartheta |\Psi(x, t)|^2 \Psi(x, t), \\ \Psi(x, 0) = \Psi_0(x), \quad (x, t) \in \Omega \times [0, T], \end{cases} \quad (2a)$$

where  $\Delta = \partial_{x_1}^2 + \dots + \partial_{x_d}^2$  denotes the Laplacian with respect to the spatial variables  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ ,  $V : \mathbb{R}^d \rightarrow \mathbb{R}$  a real-valued potential,  $\vartheta \in \mathbb{R}$  the coupling constant, and  $\Psi : \Omega \times [0, T] \subset \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{C}$  the space-time-dependent complex-valued macroscopic wave function. Assigning for a regular function  $v : \Omega \rightarrow \mathbb{C}$  the linear differential and nonlinear multiplication operators

$$\begin{aligned} (F_1(v))(x) &= c \Delta v(x), \quad c = i, \\ (F_2(v))(x) &= \bar{c} (V(x) + \vartheta |v(x)|^2) v(x), \quad \bar{c} = -i, \\ &x \in \Omega, \end{aligned} \quad (2b)$$

and setting  $u(t) = \Psi(\cdot, t)$  for  $t \in [0, T]$ , we retain the general formulation (1).

**Nonlinear parabolic equation.** By analogy to the time-dependent Gross–Pitaevskii equation (2a), we consider the parabolic problem

$$\begin{cases} \partial_t U(x, t) = \Delta U(x, t) + V(x) U(x, t) + \vartheta |U(x, t)|^2 U(x, t), \\ U(x, 0) = U_0(x), \quad (x, t) \in \Omega \times [0, T], \end{cases} \quad (3a)$$

for a real-valued solution  $U : \Omega \times [0, T] \subset \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ . Accordingly, it

corresponds to (2b) with different constant

$$\begin{aligned} (F_1(v))(x) &= c \Delta v(x), \quad c = 1, \\ (F_2(v))(x) &= \bar{c} (V(x) + \vartheta |v(x)|^2) v(x), \quad \bar{c} = 1, \\ x &\in \Omega, \end{aligned} \tag{3b}$$

and, setting  $u(t) = U(\cdot, t)$  for  $t \in [0, T]$ , we obtain again the general form (1). It is noteworthy that the parabolic equation (3a) arises in ground state and excited state computations, see for instance [5,14].

**Splitting methods.** Within multiple scopes, for ordinary differential equations and time-dependent partial differential equations, for linear problems as well as nonlinear problems, a variety of works has confirmed the benefits of operator splitting methods regarding desirable features that are subsumed under the central concepts stability, efficiency, and preservation of conserved quantities. For general information, we refer to [21,27]. Specific studies in the context of Schrödinger equations are given, e.g., in [6,8,10,32]. In order to make our subsequent survey of standard and modified operator splitting methods for nonlinear evolution equations accessible to a broader readership, we do not presume the knowledge of the formal calculus of Lie-derivatives and explain the required elementary means on occasion, see [33] for a detailed exposition.

**Linear case.** As an illustrative example, we state the simplest representative of standard splitting methods, the first-order Lie–Trotter splitting method, for a system of linear differential equations

$$\begin{cases} \frac{d}{dt} u(t) = A u(t) + B u(t), \\ u(0) = u_0, \quad t \in [0, T], \end{cases} \tag{4}$$

defined by non-commuting time-independent complex matrices  $A, B \in \mathbb{C}^{M \times M}$ . For a positive integer number  $N \in \mathbb{N}$  with associated time increment and equidistant grid points

$$\tau = \frac{T}{N}, \quad t_n = n\tau, \quad n \in \{0, 1, \dots, N\},$$

numerical approximations to the exact solution values are obtained by the recurrence

$$u_{n+1} = e^{\tau B} e^{\tau A} u_n \approx u(t_{n+1}), \quad n \in \{0, 1, \dots, N-1\}.$$

Higher-order splitting methods for (4) involve the action of several matrix exponentials on the current approximation and can be cast into the format

$$u_{n+1} = e^{b_s \tau B} e^{a_s \tau A} \dots e^{b_1 \tau B} e^{a_1 \tau A} u_n \approx u(t_{n+1}), \quad n \in \{0, 1, \dots, N-1\},$$

with real or complex coefficients  $(a_j, b_j)_{j=1}^s$ , respectively.

**Nonlinear case.** Their generalisation to nonlinear evolution equations (1) is based on the composition of the solutions to the subproblems defined by  $F_1$  and  $F_2$ . We henceforth employ the compact notation

$$\begin{aligned}\frac{d}{dt} u_1(t) &= \alpha F_1(u_1(t)), & \mathcal{E}_{\tau, \alpha F_1}(u_1(t_n)) &= u_1(t_n + \tau), \\ \frac{d}{dt} u_2(t) &= \beta F_2(u_2(t)), & \mathcal{E}_{\tau, \beta F_2}(u_2(t_n)) &= u_2(t_n + \tau), \\ \alpha, \beta &\in \mathbb{C}, & t &\in [t_n, t_n + \tau],\end{aligned}\tag{5a}$$

so that a higher-order splitting method applied to (1) reads as

$$\begin{aligned}u_{n+1} &= \left( \mathcal{E}_{\tau, b_s F_2} \circ \mathcal{E}_{\tau, a_s F_1} \circ \cdots \circ \mathcal{E}_{\tau, b_1 F_2} \circ \mathcal{E}_{\tau, a_1 F_1} \right)(u_n) \approx u(t_{n+1}), \\ n &\in \{0, 1, \dots, N-1\}.\end{aligned}\tag{5b}$$

**Schemes.** In view of numerical comparisons, we introduce the coefficients of the first-order Lie–Trotter splitting method

$$s = 1, \quad a_1 = 1, \quad b_1 = 1,\tag{6a}$$

and the second-order Strang splitting method

$$s = 2, \quad a_1 = 0, \quad a_2 = 1, \quad b_1 = \frac{1}{2} = b_2.\tag{6b}$$

Setting  $s = 4$  and imposing certain order and symmetry conditions, which reduce to a cubic equation for one of the coefficients

$$48 b_2^3 - 24 b_2^2 + 1 = 0,$$

the well-known fourth-order splitting method by YOSHIDA [34]

$$\begin{aligned}s = 4, \quad a_1 = 0, \quad a_2 = 1 - 2b_2 = a_4, \quad a_3 = 4b_2 - 1, \\ b_1 = \frac{1}{2} - b_2 = b_4, \quad b_2 = \frac{1}{6} \left( 1 - \sqrt[3]{2} - \frac{1}{2} \sqrt[3]{4} \right) = b_3,\end{aligned}\tag{6c}$$

a corresponding fourth-order splitting method with complex coefficients

$$\begin{aligned}s = 4, \quad a_1 = 0, \quad a_2 = 1 - 2b_2 = a_4, \quad a_3 = 4b_2 - 1, \\ b_1 = \frac{1}{2} - b_2 = b_4, \quad b_2 = \frac{1}{6} \left( 1 + \frac{1}{2} \sqrt[3]{2} + \frac{1}{4} \sqrt[3]{4} \right) + i \frac{\sqrt{3}}{12} \left( \frac{1}{2} \sqrt[3]{4} - \sqrt[3]{2} \right) = b_3,\end{aligned}\tag{6d}$$

and its counterpart based on the complex conjugated coefficient  $\overline{b_2}$  are retained.

**Stability issues.** In connection with dissipative systems and evolution equations of parabolic type, operator splitting methods are subject to additional stability constraints. In order to explain this matter of fact, we recall that the application of a splitting method with real coefficients  $(a_j, b_j)_{j=1}^s$  to the

parabolic equation (3) involves the subproblems

$$\partial_t U(x, t) = a_j \Delta U(x, t), \quad (x, t) \in \Omega \times [t_n, t_n + \tau], \quad j \in \{1, \dots, s\}.$$

Evidently, requiring well-posedness of these subproblems or stability of the resulting splitting method, respectively, implies

$$a_j \geq 0, \quad j \in \{1, \dots, s\}.$$

This positivity condition, however, excludes higher-order schemes, since any splitting method that exceeds a second-order barrier necessarily comprises negative coefficients, see for example [7,28,29]. Specifically, this holds true for the fourth-order splitting method by YOSHIDA

$$a_3 \approx -1.7 < 0,$$

see (6c). A feasible remedy to this issue is the design of splitting methods with complex coefficients  $(a_j, b_j)_{j=1}^s$  such that

$$\Re(a_j) \geq 0, \quad j \in \{1, \dots, s\},$$

see, e.g., [11,17]. The fourth-order scheme (6d) indeed fulfills these constraints

$$a_1 = 0, \quad \Re(a_2) = \Re(a_4) \approx 0.3 > 0, \quad \Re(a_3) \approx 0.4 > 0.$$

For further considerations in the context of the imaginary time propagation of the linear Schrödinger equation by complex splitting methods, we refer to [4].

**Modified potential splitting methods for linear equations.** Reviving former work by RUTH and SUZUKI, see for instance [26,29,30], a favourable alternative to standard operator splitting methods was proposed by CHIN. In a seminal work [12], he developed a famous fourth-order scheme of splitting type that comprises positive coefficients and hence overcomes the second-order barrier for standard splitting methods with real coefficients. Expressed in his own words, the basic idea is to incorporate *an additional higher order composite operator* so that the implementation of *one algorithm requires only one evaluation of the force and one evaluation of the force and its gradient*. For linear evolution equations (4), the resulting scheme takes the form

$$u_{n+1} = e^{\frac{1}{6}\tau B} e^{\frac{1}{2}\tau A} e^{\frac{2}{3}\tau B - \frac{1}{72}\tau^3[B, [B, A]]} e^{\frac{1}{2}\tau A} e^{\frac{1}{6}\tau B} u_n \approx u(t_{n+1}), \quad (7)$$

$$n \in \{0, 1, \dots, N-1\}.$$

Here, the iterated commutator of complex matrices  $A, B \in \mathbb{C}^{M \times M}$  is given by

$$[B, A] = BA - AB, \quad [B, [B, A]] = B^2A - 2BAB + AB^2. \quad (8a)$$

More generally, for linear differential and multiplication operators

$$\begin{aligned} (A v)(x) &= c \Delta v(x), \\ (B v)(x) &= \bar{c} V(x) v(x), \\ x &\in \Omega, \quad c \in \mathbb{C}, \end{aligned} \tag{8b}$$

retained from (2) and (3) for the special choice  $\vartheta = 0$ , a straightforward calculation yields a linear multiplication operator that depends on the Euclidean norm of the gradient of the potential

$$\begin{aligned} &[B, [B, A]]v(x) \\ &= \bar{c} |c|^2 \left( (V(x))^2 \Delta v(x) - 2V(x) \Delta(V(x) v(x)) + \Delta\left((V(x))^2 v(x)\right) \right) \tag{8c} \\ &= 2\bar{c} |c|^2 (\nabla V(x))^T \nabla V(x) v(x), \quad x \in \Omega. \end{aligned}$$

This explains the common notion *force-gradient operator splitting method* or *modified potential operator splitting method* for the scheme (7) and related splitting methods in the context of classical or quantum many-body problems and beyond. More recent contributions that exploit (7) for linear ordinary and partial differential equations are, e.g., [2,13,24,25].

**Objectives and outline.** The main objective of the present work is the extension of the modified potential operator splitting method (7) to nonlinear evolution equations of parabolic and Schrödinger type. So far, this scheme has been introduced and studied merely for linear evolution equations. In Sections 2 and 3, we state the formal generalisation to the nonlinear case and then substantiate it for the Gross–Pitaevskii equation (2) and its parabolic analogue (3). A fundamental invariance principle that includes a known result for standard splitting methods as a special case is deduced in Section 4. In Section 5, we detail the implementation of the novel modified operator splitting method based on a Fourier spectral space discretisation and provide numerical comparisons with standard splitting methods. The observed order reduction of Yoshida’s fourth-order complex splitting method (6d) is analysed in the appendix.

## 2 Modified operator splitting method

**Formal generalisation.** On the basis of the formal calculus of Lie-derivatives, our educated guess to generalise the modified potential operator splitting method (7) to the significantly more involved case of a nonlinear

evolution equation (1) is

$$u_{n+1} = \left( \mathcal{E}_{\tau, \frac{1}{6}F_2} \circ \mathcal{E}_{\tau, \frac{1}{2}F_1} \circ \mathcal{E}_{\tau, \frac{2}{3}F_2 - \frac{1}{72}\tau^2 G_2} \circ \mathcal{E}_{\tau, \frac{1}{2}F_1} \circ \mathcal{E}_{\tau, \frac{1}{6}F_2} \right) u_n \approx u(t_{n+1}), \quad (9a)$$

$$n \in \{0, 1, \dots, N-1\}.$$

Presuming well-definedness on a suitably chosen domain, the nonlinear operator  $G_2$  replacing the iterated commutator is given by

$$\begin{aligned} G_1(v) &= F_2'(v) F_1(v) - F_1'(v) F_2(v), \\ G_2(v) &= F_2'(v) G_1(v) - G_1'(v) F_2(v). \end{aligned} \quad (9b)$$

We here employ a simplified notation for the arising derivatives, which are determined as Gâteaux derivatives generalising directional derivatives

$$H'(v)w = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \left( H(v + \varepsilon w) - H(v) \right), \quad (10)$$

see also [18]. According to (5), the decisive operator is associated with a nonlinear evolution equation that comprises the time increment as parameter

$$\begin{aligned} \frac{d}{dt} \tilde{u}_2(t) &= \beta_1 F_2(\tilde{u}_2(t)) + \beta_2 \tau^2 G_2(\tilde{u}_2(t)), \quad t \in [t_n, t_n + \tau], \\ \mathcal{E}_{\tau, \beta_1 F_2 + \beta_2 \tau^2 G_2}(\tilde{u}_2(t_n)) &= \tilde{u}_2(t_n + \tau), \quad \beta_1, \beta_2 \in \mathbb{R}. \end{aligned} \quad (11)$$

**Specification and implementation.** In the subsequent sections, we complete the remaining tasks. We first specify the iterated commutator (9b) for the Gross–Pitaevskii equation (2) and contrast it to the result obtained for the parabolic counterpart (3). Then, we deduce an invariance principle that has a substantial impact on the efficiency of the modified operator splitting method (9) when applied to the time-dependent Gross–Pitaevskii equation (2). Implementation issues as well as strategies to reduce the computational cost for parabolic equations are finally discussed in Section 5.

### 3 Iterated commutators

Generally speaking, the appropriate framework for the extension of iterated commutators for matrices or linear operators, see (8), to nonlinear operators is provided by the formal calculus of Lie-derivatives. In the present work, we concretise and verify the heuristic characterisation (9) for relevant applications, the time-dependent Gross–Pitaevskii equation (2) and the related parabolic equation (3). In the context of Schrödinger equations, the arising functions  $v, w : \Omega \rightarrow \mathbb{C}$  take complex values, whereas it suffices to consider real-valued functions  $v, w : \Omega \rightarrow \mathbb{R}$  for parabolic problems. In both cases, suitable regularity requirements apply. For notational simplicity, we omit the dependence on the spatial variable.

**Derivatives.** The Gâteaux derivatives of the linear differential and nonlinear multiplication operators defined in (2) and (3) are given by

$$\begin{aligned} F_1(v) &= c \Delta v, \\ F_1'(v) w &= c \Delta w, \\ F_2(v) &= \bar{c} \left( V + \vartheta |v|^2 \right) v = \bar{c} \left( V v + \vartheta v^2 \bar{v} \right), \\ F_2'(v) w &= \bar{c} \left( V w + 2 \vartheta \bar{v} v w + \vartheta v^2 \bar{w} \right), \end{aligned}$$

see also (10).

**First commutators.** On the one hand, by performing differentiation twice, we obtain

$$\begin{aligned} F_1'(v) F_2(v) &= |c|^2 \Delta \left( V v + \vartheta v^2 \bar{v} \right) \\ &= |c|^2 \left( \Delta V v + 2 (\nabla V)^T \nabla v + V \Delta v \right) \\ &\quad + |c|^2 \vartheta \left( 2 \Delta v \bar{v} v + \Delta \bar{v} v^2 + 2 (\nabla v)^T \nabla v \bar{v} + 4 (\nabla v)^T \nabla \bar{v} v \right). \end{aligned}$$

On the other hand, a simple replacement yields

$$F_2'(v) F_1(v) = |c|^2 \left( V \Delta v + 2 \vartheta \bar{v} v \Delta v \right) + \bar{c}^2 \vartheta v^2 \Delta \bar{v}.$$

As a consequence, the difference is given by

$$\begin{aligned} G_1(v) &= F_2'(v) F_1(v) - F_1'(v) F_2(v) \\ &= -|c|^2 \left( \Delta V v + 2 (\nabla V)^T \nabla v \right) \\ &\quad + \left( \bar{c}^2 - |c|^2 \right) \vartheta \Delta \bar{v} v^2 - 2 |c|^2 \vartheta \left( (\nabla v)^T \nabla v \bar{v} + 2 (\nabla v)^T \nabla \bar{v} v \right). \end{aligned}$$

For the parabolic case (3) with  $c = 1$  this implies

$$G_1(v) = -\Delta V v - 2 (\nabla V)^T \nabla v - 6 \vartheta (\nabla v)^T \nabla v v,$$

and the analogous result for the Schrödinger case (2) with  $c = i$  is

$$G_1(v) = -\Delta V v - 2 (\nabla V)^T \nabla v - 2 \vartheta \left( \Delta \bar{v} v^2 + (\nabla v)^T \nabla v \bar{v} + 2 (\nabla v)^T \nabla \bar{v} v \right).$$

**Iterated commutators.** The iterated commutator associated with the parabolic equation (3) results from straightforward but lengthy calculations

$$\begin{aligned} G_2(v) &= 2 \left( (\nabla V)^T (\nabla V) + \vartheta \tilde{G}_2(v) \right) v, \\ \tilde{G}_2(v) &= -\Delta V v^2 + 6 (\nabla V)^T (\nabla v) v + 6 \left( V + 2 \vartheta v^2 \right) (\nabla v)^T (\nabla v), \end{aligned} \tag{12}$$



and for the Gross–Pitaevskii equation (2), we instead arrive at

$$\begin{aligned} G_2(v) &= -2i \left( (\nabla V)^T (\nabla V) - 2\vartheta \left( \tilde{G}_{21}(v) + \vartheta \tilde{G}_{22}(v) \right) \right) v, \\ \tilde{G}_{21}(v) &= |v|^2 \Delta V, \\ \tilde{G}_{22}(v) &= |v|^2 \left( 2 \Re(\bar{v} \Delta v) + 3 (\nabla \bar{v})^T (\nabla v) \right) + \Re(\bar{v}^2 (\nabla v)^T (\nabla v)). \end{aligned} \quad (13)$$

For the special case of linear multiplication operators, i.e.  $\vartheta = 0$ , we indeed recover (8). It is also noteworthy that the operator in (13) involves  $\Delta v$  and hence implies stronger regularity requirements on  $v$  compared to the operator in (12), which only comprises the gradient  $\nabla v$ .

#### 4 Invariance principle

In this section, we establish a fundamental result for the evolution operator

$$\mathcal{E}_{\tau, \beta_1 F_2 + \beta_2 \tau^2 G_2}, \quad \tau, \beta_1, \beta_2 \in \mathbb{R},$$

that has important implications concerning the efficient implementation of the modified operator splitting method (9) for nonlinear Schrödinger equations such as (2), see also (11) and (13). Furthermore, it is connected with the significance of the modified operator splitting method as a geometric integrator, see [16,19] and references given therein. We point out that the invariance principle, presented and applied in the context of (9), is a substantial extension of the known result for standard splitting methods, retained for the special choice  $\beta_2 = 0$ .

**Notation.** With regard to a compact formulation as abstract evolution equation, we again omit the dependence of the potential and a regular complex-valued function  $v : \Omega \rightarrow \mathbb{C}$  on the spatial variable. Besides, for accomplishing relations of the form

$$F_2(v) = \bar{c} f_1(v) v, \quad G_2(v) = \bar{c} f_2(v) v,$$

it is convenient to introduce the abbreviations

$$\begin{aligned} f(v) &= \beta_1 f_1(v) + \beta_2 \tau^2 f_2(v), \\ f_1(v) &= V + \vartheta g_1(v), \quad f_2(v) = 2 (\nabla V)^T (\nabla V) - 4 \vartheta g_6(v), \\ g_1(v) &= |v|^2, \quad g_2(v) = \Re(\bar{v} \Delta v), \\ g_3(v) &= (\nabla \bar{v})^T (\nabla v), \quad g_4(v) = \Re(\bar{v}^2 (\nabla v)^T (\nabla v)), \\ g_5(v) &= \vartheta \left( 2 g_2(v) + 3 g_3(v) \right), \quad g_6(v) = g_1(v) \left( \Delta V + g_5(v) \right) + \vartheta g_4(v). \end{aligned} \quad (14)$$

**Theorem (Invariance principle).** The solution to the subproblem

$$\begin{cases} \frac{d}{dt} \psi(t) = -i f(\psi(t)) \psi(t), \\ \psi(0) = \psi_0, \quad t \in [0, \tau], \end{cases}$$

with defining function introduced in (14) satisfies the invariance principle

$$f(\psi(t)) = f(\psi_0), \quad t \in [0, \tau].$$

**Proof.** In order to demonstrate that the invariance principle holds, we determine the Gâteaux derivatives of the defining functions

$$\begin{aligned} f'(v) w &= \beta_1 f'_1(v) w + \beta_2 \tau^2 f'_2(v) w, \\ f'_1(v) w &= \vartheta g'_1(v) w, \quad f'_2(v) w = -4 \vartheta g'_6(v) w, \\ g'_1(v) w &= 2 \Re(\bar{v} w), \quad g'_2(v) w = \Re(\Delta \bar{v} w) + \Re(\bar{v} \Delta w), \\ g'_3(v) w &= 2 \Re((\nabla \bar{v})^T (\nabla w)), \\ g'_4(v) w &= 2 \Re(\bar{v} \bar{w} (\nabla v)^T (\nabla v)) + 2 \Re(\bar{v}^2 (\nabla v)^T (\nabla w)), \\ g'_5(v) w &= 2 \vartheta g'_2(v) w + 3 \vartheta g'_3(v) w, \\ g'_6(v) w &= (\Delta V + g_5(v)) g'_1(v) w + g_1(v) g'_5(v) w + \vartheta g'_4(v) w, \end{aligned}$$

where suitable regularity requirements apply to  $v, w : \Omega \rightarrow \mathbb{C}$ . Observing that the potential  $V$  and the basic components  $g_1, g_2, g_3, g_4$  define real-valued functions, we have

$$\begin{aligned} (g_j(v))(x) &\in \mathbb{R}, \quad j \in \{1, \dots, 6\}, \\ (f_k(v))(x) &\in \mathbb{R}, \quad k \in \{1, 2\}, \quad (f(v))(x) \in \mathbb{R}, \quad x \in \Omega. \end{aligned}$$

Evidently, this implies that the following composition vanishes

$$g'_1(v) (i f(v) v) = 2 \Re(i |v|^2 f(v)) = 0.$$

Certain contributions originating from the iterated commutator, however, re-

quire a closer examination, namely

$$\begin{aligned}
& g'_2(v) \left( \mathfrak{i} f(v) v \right) \\
&= \Re \left( \mathfrak{i} f(v) \Re(\bar{v} \Delta v) \right) + \Re(\mathfrak{i} |v|^2 \Delta f(v)) + 2 \Re \left( \mathfrak{i} \bar{v} (\nabla f(v))^T \nabla v \right) \\
&= 2 \Re \left( \mathfrak{i} \bar{v} (\nabla f(v))^T \nabla v \right), \\
& g'_3(v) \left( \mathfrak{i} f(v) v \right) \\
&= 2 \Re \left( \mathfrak{i} v (\nabla f(v))^T \nabla \bar{v} \right) + 2 \Re \left( \mathfrak{i} f(v) (\nabla \bar{v})^T \nabla v \right) \\
&= 2 \Re \left( \mathfrak{i} v (\nabla f(v))^T \nabla \bar{v} \right), \\
& g'_4(v) \left( \mathfrak{i} f(v) v \right) \\
&= -2 \Re \left( \mathfrak{i} f(v) \bar{v}^2 (\nabla v)^T \nabla v \right) + 2 \Re \left( \mathfrak{i} |v|^2 \bar{v} (\nabla f(v))^T \nabla v \right) \\
&\quad + 2 \Re \left( \mathfrak{i} f(v) \bar{v}^2 (\nabla v)^T \nabla v \right) \\
&= 2 \Re \left( \mathfrak{i} |v|^2 \bar{v} (\nabla f(v))^T \nabla v \right).
\end{aligned}$$

On the basis of these identities, we conclude

$$\begin{aligned}
& \left( g_1(v) \left( 2 g'_2(v) + 3 g'_3(v) \right) + g'_4(v) \right) \left( \mathfrak{i} f(v) v \right) \\
&= 12 \Re \left( \mathfrak{i} |v|^2 (\nabla f(v))^T \Re(\bar{v} \nabla v) \right) = 0.
\end{aligned}$$

This proves that any composition of the special form

$$\begin{aligned}
& f'(v) \left( \mathfrak{i} f(v) v \right) \\
&= \vartheta \left( \beta_1 - 4 \beta_2 \tau^2 (\Delta V + g_5(v)) \right) g'_1(v) \left( \mathfrak{i} f(v) v \right) \\
&\quad - 4 \vartheta^2 \beta_2 \tau^2 \left( g_1(v) \left( 2 g'_2(v) + 3 g'_3(v) \right) + g'_4(v) \right) \left( \mathfrak{i} f(v) v \right) \\
&= 0
\end{aligned}$$

vanishes. As a consequence, the time-derivative of the decisive function is equal to zero

$$\frac{d}{dt} f(\psi(t)) = f'(\psi(t)) \frac{d}{dt} \psi(t) = -f'(\psi(t)) \left( \mathfrak{i} f(\psi(t)) \psi(t) \right) = 0, \quad t \in [0, \tau],$$

and hence, the desired identity follows

$$f(\psi(t)) = f(\psi_0), \quad t \in [0, \tau].$$

**Summary.** For later use, we summarise the above considerations for the

Gross–Pitaevskii equation (2). The realisation of the modified operator splitting method

$$\begin{aligned} \psi_{n+1} &= \left( \mathcal{E}_{\tau, \frac{1}{6}F_2} \circ \mathcal{E}_{\tau, \frac{1}{2}F_1} \circ \mathcal{E}_{\tau, \frac{2}{3}F_2 - \frac{1}{72}\tau^2 G_2} \circ \mathcal{E}_{\tau, \frac{1}{2}F_1} \circ \mathcal{E}_{\tau, \frac{1}{6}F_2} \right) \psi_n, \\ n &\in \{0, 1, \dots, N-1\}, \end{aligned}$$

results in the numerical integration of the linear Schrödinger equation

$$\frac{d}{dt} \psi(t) = i \Delta \psi(t), \quad t \in [t_n, t_n + \frac{\tau}{2}], \quad \mathcal{E}_{\tau, \frac{1}{2}F_1}(\psi(t_n)) = \psi(t_n + \frac{\tau}{2}),$$

and the evaluation of the solution representation

$$\mathcal{E}_{\tau, \beta_1 F_2 + \beta_2 \tau^2 G_2}(\psi_0) = e^{-i\tau(\beta_1 f_1(\psi_0) + \beta_2 \tau^2 f_2(\psi_0))} \psi_0, \quad \tau, \beta_1, \beta_2 \in \mathbb{R}, \quad (15)$$

which holds pointwise for any argument  $x \in \Omega$ , see (14) for the definitions of  $f_1$  and  $f_2$ .

## 5 Numerical results

In the following, we illustrate the stability and global error behaviour of the novel modified operator splitting method (9) for the time-dependent Gross–Pitaevskii equation (2) and its parabolic analogue (3). The numerical tests, performed in one, two, and three space dimensions, in particular confirm the theoretical considerations of Sections 2 to 4. For the purpose of comparison, we in addition include the corresponding results for widely-used standard splitting methods.

**Implementation.** The practical realisation of standard operator splitting methods such as (6) and of the modified operator splitting method (9), respectively, requires the time integration of the subproblems involving the Laplacian and the nonlinear multiplication operator. In our implementation, we make use of fast Fourier techniques, which are based on the following considerations.

- (i) *Space grid.* With regard to the employed Fourier spectral space discretisation, we replace the underlying unbounded domain by a Cartesian product of sufficiently large intervals

$$a = 10, \quad x \in [-a, a]^d \subset \Omega,$$

and choose the total number of equidistant spatial grid points according to the dimension

$$d = 1 : M = 512, \quad d = 2 : M = 128^2, \quad d = 3 : M = 64^3.$$

- (ii) *Derivatives.* The iterated commutator arising in the modified operator splitting method (9) for (2) and (3) involves the gradient  $\nabla V$  and the Laplacian  $\Delta V$  of the space-dependent potential, which we may assume to be known analytically. Otherwise, we employ the approach described subsequently. The numerical computation of the spatial derivatives  $\nabla v$  and  $\Delta v$ , where  $v$  represents the current value of the time-discrete solution, is traced back to a fast Fourier transform, pointwise multiplications, and an inverse fast Fourier transform. Denoting by  $(\mathcal{F}_m)_{m \in \mathbb{Z}^d}$  the Fourier functions with periodicity domain  $[-a, a]^d$ , by  $(\mu_m)_{m \in \mathbb{Z}^d}$  the purely imaginary eigenvalues associated with the first spatial derivatives, and by  $(\lambda_m)_{m \in \mathbb{Z}^d}$  the corresponding real eigenvalues of the Laplace operator

$$\begin{aligned}\mathcal{F}_m(x) &= (2a)^{-\frac{d}{2}} e^{i\pi m_1 (\frac{x_1}{a} + 1)} \dots e^{i\pi m_d (\frac{x_d}{a} + 1)}, \\ \nabla \mathcal{F}_m &= \mu_m \mathcal{F}_m, \quad \mu_m = \frac{i\pi m}{a} \in \mathbb{C}^{d \times 1}, \\ \Delta \mathcal{F}_m &= \lambda_m \mathcal{F}_m, \quad \lambda_m = -\frac{\pi^2 |m|^2}{a^2} \in \mathbb{R}, \\ m &= (m_1, \dots, m_d) \in \mathbb{Z}^d, \quad x = (x_1, \dots, x_d) \in [-a, a]^d,\end{aligned}$$

the following formal representations hold

$$\begin{aligned}v &= \sum_{m \in \mathbb{Z}^d} v_m \mathcal{F}_m, \quad v_m = \int_{[-a, a]^d} v(x) \mathcal{F}_{-m}(x) dx, \quad m \in \mathbb{Z}^d, \\ \nabla v &= \sum_{m \in \mathbb{Z}^d} \mu_m v_m \mathcal{F}_m, \quad \Delta v = \sum_{m \in \mathbb{Z}^d} \lambda_m v_m \mathcal{F}_m.\end{aligned}$$

Their realisation relies on a suitable truncation of the infinite index set  $\mathcal{M} \subset \mathbb{Z}^d$  so that  $|\mathcal{M}| = M$  and quadrature approximations by the trapezoidal rule.

- (iii) *Linear subproblem.* Formally, the solution to the linear subproblem is given by a Fourier series

$$\begin{aligned}\frac{d}{dt} u_1(t) &= \alpha F_1(u(t)) = c \alpha \Delta u_1(t), \quad t \in [t_n, t_n + \tau], \quad \alpha \in \mathbb{R}, \\ \mathcal{E}_{\tau, \alpha F_1}(u(t_n)) &= \mathcal{E}_{\tau, \alpha F_1} \left( \sum_{m \in \mathbb{Z}^d} u_{1,m}(t_n) \mathcal{F}_m \right) = \sum_{m \in \mathbb{Z}^d} e^{c \alpha \tau \lambda_m} u_{1,m}(t_n) \mathcal{F}_m,\end{aligned}$$

see also (2), (3), and (5). Again, the application of fast Fourier techniques permits the efficient computation of approximations to the spectral coefficients and the evaluation of finite sums on the equidistant spatial grid points covering the underlying domain

$$\begin{aligned}\tilde{u}_{1,m,n} &\approx u_{1,m}(t_n), \quad m \in \mathcal{M}, \\ \sum_{m \in \mathcal{M}} e^{c \alpha \tau \lambda_m} \tilde{u}_{1,m,n} \mathcal{F}_m(x) &, \quad x \in [-a, a]^d.\end{aligned}$$

- (iv) *Nonlinear subproblem (Schrödinger equation).* In the context of the Gross–Pitaevskii equation (2), we make use of the fact that the solution to the nonlinear subproblem (11) satisfies the invariance principle

deduced in Section 4. Consequently, it simply remains to evaluate the representation (15) on the equidistant grid.

- (v) *Nonlinear subproblem (Parabolic equation).* In the case of the parabolic equation (3), we additionally apply an explicit Runge–Kutta method of order four. Due to the stiffness of the problem, the time stepsize has to be adjusted to the spatial grid width to ensure stability. Alternative approaches with improved stability properties and reduced computational costs are detailed below.

**Numerical results.** In our numerical tests, we perform the time integration of the Gross–Pitaevskii equation (2) and the parabolic equation (3), opposing the nonlinear case with  $\vartheta = 1$  to the simplified linear case with  $\vartheta = 0$ . We prescribe the Gaussian-shaped initial state

$$u_0(x) = e^{-\frac{1}{2}(x_1^2 + \dots + x_d^2)}, \quad x = (x_1, \dots, x_d) \in \mathbb{R}^d,$$

as well as the two polynomial potentials

$$V(x) = C_0 C_q \sum_{j=1}^d x_j^q,$$

$$C_2 = 1, \quad C_4 = \frac{1}{24}, \quad q \in \{2, 4\}, \quad x = (x_1, \dots, x_d) \in \mathbb{R}^d,$$

for which the needed first- and second-order derivatives are known analytically. For the special case of a quadratic potential with prefactor  $C_0 \in \{1, -1\}$  chosen accordingly to the type of the equation and  $\vartheta = 0$ , the knowledge of the exact solution

$$u(x, t) = e^{d\bar{c}t} u_0(x), \quad (x, t) \in \mathbb{R}^d \times [0, T],$$

permits the validation of the Fourier spectral space discretisation and the time-splitting approach. In the general case, we instead determine a numerical reference solution based on a refined time stepsize. The global errors of the modified operator splitting method (9) at final time  $T = 1$ , measured in a discrete  $L_2$ -norm, are compared to those obtained by the standard Lie–Trotter, Strang, and Yoshida splitting methods of non-stiff orders one, two, and four, see also (6). The obtained results, displayed in Figures 1–4, confirm the favourable behaviour of the modified operator splitting method.

**Computational cost.** In general, an expedient measure for the computational cost of the modified operator splitting method (9) for the Gross–Pitaevskii equation (2) and the parabolic equation (3), respectively, is the number of fast Fourier transforms and their inverses. Evidently, the numerical solution of two linear subproblems per time step amounts to two fast Fourier transforms and two inverse fast Fourier transforms. Besides, the time integration of the nonlinear subproblem requires the computation of space derivatives via Fourier transforms.

- (i) For nonlinear Schrödinger equations such as (2), the validity of the invariance property permits to significantly reduce the cost related to the evaluation of  $\mathcal{E}_{\tau, \frac{2}{3}F_2 - \frac{1}{72}\tau^2 G_2}$ . Due to the fact that the spectral coefficients of the current numerical solution  $v$  are available, the computation of gradient  $\nabla v$  and Laplacian  $\Delta v$  results in  $d+1$  inverse fast Fourier transforms.
- (ii) For the parabolic equation (3), a favourable approach is based on the following considerations. The presence of the additional factor  $\tau^2$  in connection with the double commutator permits to use an approximation by means of the the second-order Strang splitting method, that is

$$\mathcal{E}_{\tau, \frac{2}{3}F_2 - \frac{1}{72}\tau^2 G_2} \approx \mathcal{E}_{\frac{1}{2}\tau, \frac{2}{3}F_2} \circ \mathcal{E}_{\tau, -\frac{1}{72}\tau^2 G_2} \circ \mathcal{E}_{\frac{1}{2}\tau, \frac{2}{3}F_2}.$$

On the one hand, an explicit representation of the evolution operator associated with the nonlinear ordinary differential equation

$$\frac{d}{dt} u(x, t) = \left( V(x) + \vartheta |u(x, t)|^2 \right) u(x, t), \quad x \in \Omega, \quad t \in [0, \tau],$$

is known. On the other hand, for the time integration of the nonlinear subproblem involving the double commutator, it suffices to apply the first-order explicit Euler method. Numerical tests confirm the enhanced stability, accuracy, and efficiency of the resulting approach, in particular, in case of a fourth-order polynomial potential, see Figure 5.

## 6 Conclusions

In the present work, we have introduced a general framework for the generalisation of CHIN's fourth-order modified potential operator splitting method (7) to nonlinear evolution equations. To the best of our knowledge, this matter is novel and of major interest from theoretical and practical perspectives.

Moreover, we have specified the resulting fourth-order modified operator splitting method (9) for the time-dependent Gross–Pitaevskii equation (2) and its parabolic counterpart (3). It seems likely that our approach and the drawn conclusions also extend to Schrödinger equations involving nonlinear and possibly nonlocal terms of the form  $g(|\Psi(x, t)|) \Psi(x, t)$ .

Due to the fact that our numerical tests have confirmed the favourable performance of the proposed fourth-order scheme in comparison with standard real and complex splitting methods of order four, it is natural to extend our considerations in various respects.

Proceeding our recent work [9] for linear evolution equations of parabolic and Schrödinger type, we find it promising to design high-order modified opera-

tor splitting methods for nonlinear evolution equations that are optimal with regard to a preselected criterium such as efficiency. In contrast to the linear case, we have to take into account the additional costs for the evaluation of second iterated commutators and that higher-order iterated commutators will not vanish, in general, see (8) as well as (12) and (13). Nonetheless, based on the successful strategies for the efficient implementation of the fourth-order scheme (9), it suggests itself to address an in-depth analysis of similarly structured modified operator splitting methods.

A desirable feature of the proposed fourth-order modified operator splitting method is the positivity of the coefficients. Concerning the design of high-order schemes, in light of contributions on the linear case, see for instance [3,13], this will necessitate further deliberations on the appropriate format. Evidently, positivity is intrinsically related to the issue of well-posedness and stability for parabolic problems. Besides, it affects aliasing effects on truncated space domains and the incorporation of artificial boundary conditions, which is of particular interest in the context of nonlinear Schrödinger equations, see for example [1] and references given therein.

Furthermore, we intend to carry out a rigorous convergence analysis of modified operator splitting methods applied to nonlinear evolution equations. We point out that the study for the linear case [20] together with the formal calculus of Lie-derivatives provides a guiding principle. However, as seen in Section 3, the accomplishment for specific nonlinear evolution equations involving unbounded operators requires careful calculations and investigations, see also [31].

## Acknowledgements

The authors dedicate this work to Arieh Iserles due to his seminal contributions in the area of numerical analysis and geometric numerical integration. This work has been supported by Ministerio de Ciencia e Innovación (Spain) through projects PID2019-104927GB-C21 and PID2019-104927GB-C22, MCIN/AEI/10.13039/501100011033, ERDF (*A way of making Europe*).

## References

- [1] X. ANTOINE, A. ARNOLD, CH. BESSE, M. EHRHARDT, A. SCHÄDLE. *A review of transparent and artificial boundary conditions techniques for linear and nonlinear Schrödinger equations*. Commun. Comput. Phys. 4 (2008) 729–796.



- [2] J. AUER, E. KROTSCHECK, S. CHIN. *A fourth-order real-space algorithm for solving local Schrödinger equations*. J. Chem. Phys. 115 (2001) 6841–6846.
- [3] W. AUZINGER, H. HOFSTÄTTER, O. KOCH. *Non-existence of generalized splitting methods with positive coefficients of order higher than four*. Appl. Math. Letters 97 (2019) 48–52.
- [4] P. BADER, S. BLANES, F. CASAS. *Solving the Schrödinger eigenvalue problem by the imaginary time propagation technique using splitting methods with complex coefficients*. J. Chem. Phys. 139 (2013) 124117.
- [5] W. BAO, Q. DU. *Computing the ground state solution of Bose–Einstein condensates by a normalized gradient flow*. SIAM J. Sci. Comput. 25 (2004) 1674–1697.
- [6] W. BAO, S. JIN, P. MARKOWICH. *On time-splitting spectral approximations for the Schrödinger equation in the semiclassical regime*. J. Comput. Phys. 175 (2002) 487–524.
- [7] S. BLANES, F. CASAS. *On the necessity of negative coefficients for operator splitting schemes of order higher than two*. Appl. Numer. Math. 54 (2005) 23–37.
- [8] S. BLANES, F. CASAS, A. MURUA. *An efficient algorithm based on splitting for the time integration of the Schrödinger equation*. J. Comput. Phys. 303 (2015) 396–412.
- [9] S. BLANES, F. CASAS, C. GONZÁLEZ, M. THALHAMMER. *Efficient splitting methods based on modified potentials: numerical integration of linear parabolic problems and imaginary time propagation of the Schrödinger equation*. To appear in Commun. Comput. Phys. (2023).
- [10] M. CALIARI, S. ZUCCHER. *A fast time splitting finite difference approach to Gross–Pitaevskii equations*. Commun. Comput. Phys. 29 (2021) 1336–1364.
- [11] F. CASTELLA, P. CHARTIER, S. DECOMBES, G. VILMART. *Splitting methods with complex times for parabolic equations*. BIT 49 (2009) 487–508.
- [12] S. CHIN. *Symplectic integrators from composite operator factorizations*. Phys. Lett. A 226 (1997) 344–348.
- [13] S. CHIN. *Structure of positive decomposition of exponential operators*. Phys. Rev. E 71 (2005) 016703.
- [14] I. DANAILA, B. PROTAS. *Computation of ground states of the Gross–Pitaevskii functional via Riemannian optimization*. SIAM J. Sci. Comput. 39 (2017) 1102–1129.
- [15] E. GROSS. *Structure of a quantized vortex in boson systems*. Nuovo Cimento 20 (1961) 454–477.
- [16] E. HAIRER, CH. LUBICH, G. WANNER. *Geometric Numerical Integration*. Springer, Second Edition, 2006.

- [17] E. HANSEN, A. OSTERMANN. *High order splitting methods for analytic semigroups exist*. BIT 49 (2009) 527–542.
- [18] E. HILLE, R. PHILLIPS. *Functional Analysis and Semi-Groups*. American Mathematical Society, 1974.
- [19] A. ISERLES. *A First Course in the Numerical Analysis of Differential Equations*. Cambridge University Press, Second Edition, 2008.
- [20] E. KIERI. *Stiff convergence of force-gradient operator splitting methods*. Appl. Numer. Math. 94 (2015) 33–45.
- [21] R. MCLACHLAN, G. QUISPÉL. *Splitting methods*. Acta Numerica 11 (2002) 341–434.
- [22] A. MESSIAH. *Quantum Mechanics*. Dover, 1999.
- [23] L. PITAEVSKII. *Vortex lines in an imperfect Bose gas*. Sov. Phys. JETP 13 (1961) 451–454.
- [24] I. OMEL'YAN, I. MRYGŁOD, R. FOLK. *On the construction of high order force gradient algorithms for integration of motion in classical and quantum systems*. Phys. Rev. E 66 (2002) 026701.
- [25] I. OMEL'YAN, I. MRYGŁOD, R. FOLK. *Symplectic analytically integrable decomposition algorithms: classification, derivation, and application to molecular dynamics, quantum and celestial mechanics simulations*. Comput. Phys. Comm. 151 (2003) 272–314.
- [26] R. RUTH. *A canonical integration technique*. IEEE Trans. Nucl. Sci. 30 (1983) 26–69.
- [27] J. SANZ-SERNA, M. CALVO. *Numerical Hamiltonian Problems*. Dover, 2018.
- [28] Q. SHENG. *Solving linear partial differential equations by exponential splitting*. IMA J. Numer. Anal. 9 (1989) 199–212.
- [29] M. SUZUKI. *General theory of fractal path integrals with applications to many-body theories and statistical physics*. J. Math. Phys. 32 (1991) 400–407.
- [30] M. SUZUKI. *Hybrid exponential product formulas for unbounded operators with possible applications to Monte Carlo simulations*. Phys. Lett. A 201 (1995) 425–428.
- [31] M. THALHAMMER. *High-order exponential operator splitting methods for time-dependent Schrödinger equations*. SIAM J. Numer. Anal. 46 (2008) 2022–2038.
- [32] M. THALHAMMER, J. ABHAU. *A numerical study of adaptive space and time discretisations for Gross–Pitaevskii equations*. J. Comput. Phys. 231 (2012) 6665–6681.
- [33] V. VARADARAJAN. *Lie Groups, Lie Algebras, and Their Representations*. Springer, 1984.
- [34] H. YOSHIDA. *Construction of higher order symplectic integrators*. Phys. Lett. A 150 (1990) 262–268.




Fig. 1. Time integration of the Gross–Pitaevskii equation (2) involving a quadratic potential by standard splitting methods and the novel modified operator splitting method. Global errors versus time stepsizes in space dimensions  $d \in \{1, 2, 3\}$ . Nonlinear ( $\vartheta = 1$ ) versus simplified linear ( $\vartheta = 0$ ) case. Due to the validity of the invariance principle, the application of an explicit Runge–Kutta method is not needed (RK0).




Fig. 2. Time integration of the parabolic problem (3) involving a quadratic potential by standard splitting methods and the novel modified operator splitting method. Global errors versus time stepsizes in space dimensions  $d \in \{1, 2, 3\}$ . Nonlinear ( $\vartheta = 1$ ) versus simplified linear ( $\vartheta = 0$ ) case. In order to resolve the nonlinear sub-problem, a fourth-order explicit Runge–Kutta method is applied (RK4). Depending on the stiffness of the equation, stability is ensured for sufficiently small time stepsizes. For a naive implementation of the Yoshida splitting method with complex coefficients, an order reduction is observed, see Appendix A.




Fig. 3. Corresponding results for the time-dependent Gross–Pitaevskii equation (2) involving a fourth-order polynomial potential.




Fig. 4. Corresponding results for the parabolic problem (3) involving a fourth-order polynomial potential.






Fig. 5. Time integration of the one-dimensional parabolic equation (3) involving a quadratic potential (left) or a fourth-order potential (right), respectively, by the modified operator splitting method (9). Global errors versus time stepsizes. The original approach is based on the application of an explicit fourth-order Runge–Kutta method for the numerical solution of the nonlinear subproblem involving the double commutator (cf.  $\mathcal{E}_{\tau, \frac{2}{3}F_2 - \frac{1}{72}\tau^2 G_2}$ ). Alternative approaches are based on the Strang splitting method (cf.  $\mathcal{E}_{\frac{1}{2}\tau, \frac{2}{3}F_2} \circ \mathcal{E}_{\tau, -\frac{1}{72}\tau^2 G_2} \circ \mathcal{E}_{\frac{1}{2}\tau, \frac{2}{3}F_2}$ ). Here, a reduced number of (inverse) fast Fourier transforms is required and an improved accuracy is observed. Furthermore, the knowledge of the exact solution to a component (cf.  $\mathcal{E}_{\frac{1}{2}\tau, \frac{2}{3}F_2}$ ) enhances the stability behaviour of the resulting time integration method for larger time increments.

## A Appendix: Order reduction of complex splitting methods

In order to explain the observed order reduction for the complex splitting method (6d), it suffices to study a nonlinear ordinary differential equation of the form (1) with  $F_1 = 0$  and  $F_2(u) = |u|^2 u$  on a single subinterval of length  $(0, \tau)$ . More precisely, we consider the nonlinear subproblems

$$\begin{cases} \frac{d}{dt} u(t) = F_2^{(1)}(u(t)) = (u(t))^3, & \frac{d}{dt} u(t) = F_2^{(2)}(u(t)) = |u(t)|^2 u(t), \\ u(0) = u_0 \in \mathbb{R}, \quad t \in (0, \tau), & u(0) = u_0 \in \mathbb{R}, \quad t \in (0, \tau), \end{cases}$$

with coinciding real-valued solutions

$$\mathcal{E}_{\tau, F_2^{(1)}}(u_0) = u(\tau) = \mathcal{E}_{\tau, F_2^{(2)}}(u_0).$$

On the one hand, using the Taylor series expansion

$$u(\tau) = u(0) + \tau u'(0) + \frac{1}{2} \tau^2 u''(0) + \frac{1}{6} \tau^3 u'''(0) + \frac{1}{24} \tau^4 u''''(0) + \mathcal{O}(\tau^5)$$

in combination with the differential equation and derivatives thereof implies

$$u(\tau) = \left(1 + \tau u_0^2 + \frac{3}{2} \tau^2 u_0^4 + \frac{5}{2} \tau^3 u_0^6 + \frac{35}{8} \tau^4 u_0^8\right) u_0 + \mathcal{O}(\tau^5).$$

As a consequence, for any real number  $b \in \mathbb{R}$ , the relation

$$b \in \mathbb{R} : \quad u(b\tau) = \left(1 + b\tau u_0^2 + \frac{3}{2} b^2 \tau^2 u_0^4 + \frac{5}{2} b^3 \tau^3 u_0^6 + \frac{35}{8} b^4 \tau^4 u_0^8\right) u_0 + \mathcal{O}(\tau^5)$$

is valid. For complex numbers, however, the more general expansion

$$\begin{aligned} b, u_0 \in \mathbb{C} : \quad \mathcal{E}_{\tau, bF_2^{(2)}}(u_0) = & \left(1 + b\tau |u_0|^2 + \left(b^2 + \frac{1}{2} |b|^2\right) \tau^2 |u_0|^4 \right. \\ & + \left(b^3 + \left(\frac{7}{6} b + \frac{1}{3} \bar{b}\right) |b|^2\right) \tau^3 |u_0|^6 \\ & + \left(b^4 + \frac{1}{24} (46 b^2 + 6 \bar{b}^2 + 29 |b|^2) |b|^2\right) \tau^4 |u_0|^8 \Big) u_0 \\ & + \mathcal{O}(\tau^5) \end{aligned}$$

is obtained by decomposing the solution and accordingly the defining function into real and imaginary parts. In order to reproduce the approximation that corresponds to the Yoshida splitting, we impose the basic symmetry and order conditions

$$b_4 = b_1, \quad b_3 = b_2, \quad b_1 + b_2 + b_3 + b_4 = 1,$$

and then perform the fourfold composition

$$u_{\text{Splitting}}(\tau) = \left(\mathcal{E}_{\tau, b_1 F_2^{(2)}} \circ \mathcal{E}_{\tau, b_2 F_2^{(2)}} \circ \mathcal{E}_{\tau, b_2 F_2^{(2)}} \circ \mathcal{E}_{\tau, b_1 F_2^{(2)}}\right)(u_0).$$



Requiring this expansion to coincide with the expansion of the exact solution

$$u_{\text{Splitting}}(\tau) - u(\tau) = C(u_0) \Im(b_1) (1 - \Re(b_1)) \tau^3 + \mathcal{O}(\tau^4)$$

leads to a condition that obviously contradicts the order conditions for splitting methods and explains the observation of local order three and global order two. Similar arguments apply to evolution equations of Schrödinger type such as the Gross–Pitaevskii equation. But, in this context, the invariance principle permits to avoid the application of a fourth-order Runge–Kutta method to the arising nonlinear subproblem

$$\frac{d}{dt} u(t) = F_2^{(2)}(u(t)) = -i |u(t)|^2 u(t), \quad t \in (0, \tau).$$