

Modified splitting methods for Gross–Pitaevskii systems modelling Bose–Einstein condensates: Time evolution and ground state computation

Mechthild Thalhammer, Gregor Thalhammer–Thurner¹

Abstract

The year 2025 marks the 100 and 30 years anniversaries of the discovery of Bose–Einstein condensation and its successful experimental realisation. Inspired by these important research achievements, a conceptually simple approach is proposed to facilitate reliable and efficient numerical simulations. The structure of the underlying systems of coupled Gross–Pitaevskii equations suggests the use of optimised high-order operator splitting methods for dynamical evolution and ground state computation. A second-order barrier, however, prevents the applicability of standard operator splitting methods for both, time evolution as well as imaginary time propagation. An innovative alternative approach accomplishes the design of novel modified operator splitting methods that remain stable under moderate smallness assumptions on the time increments. The core idea is to incorporate commutators of the defining differential and nonlinear multiplication operators, since this permits to fulfill the basic stability requirement of positive method coefficients. Further improvements with respect to convergence at the targeted precision arise from automatic adjustments of the time stepsizes by an inexpensive local error control. The presented numerical experiments confirm the favourable performance of a specific fourth-order modified operator splitting method. Amongst others, it is demonstrated that the excellent mass and energy conservation in long-term evolutions, intrinsic attributes of geometric numerical integrators for Hamiltonian systems, is maintained for a sensible variation of the time stepsizes. Moreover, the benefits of adaptive higher-order approximations in ground state computations are illustrated.

Key words: Bose–Einstein condensation, Gross–Pitaevskii systems, Nonlinear Schrödinger equations, Time evolution, Ground state computation, Imaginary time propagation, Geometric numerical integration, Modified operator splitting methods, Fast Fourier transform, Stability, Convergence, Efficiency, Adaptivity.

¹Addresses: Mechthild Thalhammer, University of Innsbruck, Department of Mathematics, 6020 Innsbruck, Austria. Gregor Thalhammer–Thurner, Medical University of Innsbruck, Institute of Biomedical Physics, 6020 Innsbruck, Austria. Websites: techmath.uibk.ac.at/mecht, www.i-med.ac.at/dpmp/bmp. Email addresses: mechthild.thalhammer@uibk.ac.at, gregor.thalhammer@i-med.ac.at.

1 Introduction

Main objective. In light of the 100 and 30 years anniversaries of the theoretical discovery of Bose–Einstein condensation [21] and its successful experimental realisation at ultracold temperatures [1], a main objective of the present work is the introduction and exemplification of a novel and conceptually simple approach, in order to facilitate the reliable and efficient numerical simulation of multi-species Bose–Einstein condensates.

Bose–Einstein condensates. The creation of Bose–Einstein condensate collective systems occurs when a gas of bosons is cooled to nanokelvin temperatures by laser technologies, since the controlled reduction of atomic motion causes a large number of microscopic particles to form a single macroscopic quantum state at lowest energy. Worldwide, various experimental physics groups devote their current research to the study and manipulation of this particular state of matter. As an instance, we highlight the *Strongly Correlated Quantum Matter Group*² at the University of Innsbruck, led by HANNS–CHRISTOPH NÄGERL, and the recent contribution [20].

Gross–Pitaevskii systems. Fundamental mathematical models that describe the dynamical evolution of the macroscopic wave functions associated with multi-species Bose–Einstein condensates are provided by systems of coupled time-dependent Gross–Pitaevskii equations [22,27]. We focus on typical instances that are characterised by partial differential equations of Schrödinger-type comprising the Laplacian with respect to the spatial coordinates, real-valued space-dependent functions, and non-analytic nonlinearities. The defining operators reflect kinetic energies, external trapping and lattice potentials, and the interactions between the particles of different species. The physically most relevant setting corresponds to three space dimensions and cubic nonlinearities. In addition, we study the special cases of one and two dimensions with the advantages of accelerating numerical tests and simplifying the visualisations of solution profiles.

Operator splitting approach. The structures of the defining linear differential and nonlinear multiplication operators suggest the use of standard operator splitting methods combined with fast Fourier techniques for spatial discretisation. The main idea is to divide the original problem into subproblems and to apply adjusted solvers to each of the subproblems. Employing a usual time-stepping approach, the respective approximations are composed several times, whereby the time increments are scaled by certain method coefficients. The particular choice of the method coefficients, linked to the number of compositions per time step, determines the overall quantitative and qualita-

² See quantummatter.at.

tive properties of a specific splitting scheme. Detailed information on splitting methods is found in [10,26]. A small selection of contributions in the context of linear and nonlinear Schrödinger equations is [5,9,16,32–34], see also the references given therein.

Geometric numerical integrators. Optimised high-order splitting methods exhibit a favourable behaviour regarding stability, accuracy, and efficiency. By reason of their distinguishing attributes of geometric numerical integrators for Hamiltonian systems [8,23,24,31], they also show an excellent preservation of conserved quantities such as mass and energy over long times.

Limitations for non-reversible systems. However, the unified applicability of standard operator splitting methods for both, the time evolution and the imaginary time propagation of Gross–Pitaevskii systems, is limited by a second-order barrier for systems that are non-reversible in time. In other words, due to the fact that higher-order splitting methods necessarily involve negative coefficients [7] and thus suffer from severe instabilities, they are excluded from ground and excited state computations. Hence, the symmetric second-order Strang splitting method is left as canonical choice.

Modified operator splitting approach. In this contribution, we exploit an alternative approach, which we refer to as modified operator splitting, and illustrate its intrinsic benefits over standard splitting methods and other classes of composition methods, see Section 4 below. A key observation is that the incorporation of double commutators of the defining differential and multiplication operators in the nonlinear subproblems allows for the design of fourth-order schemes with positive coefficients. Within the context of *linear* differential equations, the basic concept is known as modified potential or force-gradient operator splitting, see [3,11,17,18,25,29]. In our recent work [12], we could accomplish the generalisation to a class of nonlinear partial differential equations by the formal calculus of Lie-derivatives [36]. In what follows, we establish modified operator splitting methods for the first time for Gross–Pitaevskii systems. On the one hand, we verify conservation of mass and hence stability by means of a fundamental invariance principle. On the other hand, we demonstrate that schemes with positive coefficients remain stable for their parabolic-type counterparts arising in the imaginary time propagation for ground and excited state computations, see [6,11,19,28], provided that the time stepsizes are moderately small. As a result of these well-founded considerations, sustained by numerical experiments presented in Section 5, we conclude that modified operator splitting methods with positive coefficients pave the way for reliable ground state computations and efficient time evolution. Such scenarios, the preparation of ground state solutions and subsequent dynamical manipulation, are of particular relevance in actual laboratory setups [20,30,35]. The approach and the resulting implementation are instantaneously adaptable to any other choice of the scheme, since it suffices to

redefine the method coefficients in a single component of the code.

Adaptivity in time. With regard to the evident advantages of adaptivity in time, we propose two strategies for an inexpensive local error control. According to the requirements of the considered problem and the ranges of the decisive physical parameters, these options automatically adapt the time increments to ensure stability and also enhance the targeted precision. We measure the effort of the resulting adaptive modified splitting methods by the numbers of fast Fourier transforms and their inverses, since these procedures constitute the computationally most costly components.

Efficient realisation. The validity of an invariance principle has a significant impact on the efficient realisation of modified splitting methods for time-dependent Gross–Pitaevskii systems. In essence, it justifies the replacement of nonlinear terms depending on the time-dependent solutions by expressions involving only the initial solution values. As a consequence, for the nonlinear subproblems that result from omitting the Laplacians, this invariance property provides the knowledge of the exact solutions on each of the temporal subintervals.

Numerical experiments. Our numerical experiments for model problems in one, two, and three space dimensions illustrate the benefits of a specific adaptive fourth-order modified operator splitting method in comparison with standard splitting methods based on uniform time grids. Amongst others, we confirm the favourable energy conservation in long-term evolutions and demonstrate the computation of ground states with high accuracy.

Notation. In the following, we consider suitably chosen subsets of the d -dimensional Euclidean space as underlying space domains and set

$$|x| = \sqrt{x_1^2 + \cdots + x_d^2}, \quad x = (x_1, \dots, x_d) \in \Omega \subseteq \mathbb{R}^d, \quad d \in \{1, 2, 3\}.$$

The Lebesgue space $L^2(\Omega, \mathbb{C})$ of square integrable complex-valued functions forms a Hilbert space with inner product and associated norm is defined by

$$(\varphi | \psi)_{L^2} = \int_{\Omega} \varphi(x) \overline{\psi(x)} \, dx, \quad \|\psi\|_{L^2} = \sqrt{(\psi | \psi)_{L^2}}, \quad \varphi, \psi \in L^2(\Omega, \mathbb{C}). \quad (1a)$$

Accordingly, for vector-valued functions, we denote

$$\begin{aligned} (\varphi | \psi)_{(L^2)^d} &= \sum_{j=1}^d (\varphi_j | \psi_j)_{L^2}, \quad \|\psi\|_{(L^2)^d} = \sqrt{\sum_{j=1}^d \|\psi_j\|_{L^2}^2}, \\ \varphi &= (\varphi_1, \dots, \varphi_d) \in L^2(\Omega, \mathbb{C}^d), \quad \psi = (\psi_1, \dots, \psi_d) \in L^2(\Omega, \mathbb{C}^d). \end{aligned} \quad (1b)$$

The arising derivatives of functionals are determined as Gâteaux derivatives

generalising directional derivatives

$$F'(v)w = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} (F(v + \varepsilon w) - F(v)). \quad (2)$$

Throughout, we tacitly assume that for the situations under consideration sufficiently regular solutions are ensured such that the application of high-order space and time discretisation methods is indeed expedient.

Outline. The present work is organised as follows. In Sections 2 and 3, we state Gross–Pitaevskii equations as well as extended systems modelling multi-species Bose–Einstein condensates. A compact symbolic notation permits to emphasise the common structures of the initial value problems for dynamical evolution and ground state computation. In Section 4, we first give a brief summary of standard operator splitting methods and then detail our novel approach of modified operator splitting for the time evolution and imaginary time propagation of Gross–Pitaevskii systems. We clarify the validity of an invariance principle that has a significant impact on the efficient realisation for time-dependent Gross–Pitaevskii systems and include information on the practical implementation of modified operator splitting methods based on a Fourier spectral space discretisation. Moreover, we contrast modified operator splitting methods to an alternative class of methods, operator splitting methods with complex coefficients. Comparative numerical experiments regarding the stability and accuracy behaviour of modified operator splitting methods in short-term as well as long-term evolutions for two-component systems are presented in Section 5. Concluding remarks are given in the final section 6.

2 Gross–Pitaevskii equations

Model problem. As a basic model for the nonlinear dynamics of a single-species Bose–Einstein condensate, we consider the initial value problem for a time-dependent Gross–Pitaevskii equation. For our purposes, it suffices to study the dimensionless formulation

$$\begin{cases} i \partial_t \Psi(x, t) = \Delta_\alpha \Psi(x, t) + V_{\beta, \gamma, \delta}(x) \Psi(x, t) + \vartheta |\Psi(x, t)|^2 \Psi(x, t), \\ \Psi(x, t_0) = \Psi_0(x), \quad (x, t) \in \Omega \times [t_0, T]. \end{cases} \quad (3a)$$

Here, we denote by $i \in \mathbb{C}$ the imaginary unit, by

$$\Psi : \Omega \times [t_0, T] \longrightarrow \mathbb{C} : (x, t) \longmapsto \Psi(x, t) \quad (3b)$$

the complex-valued macroscopic wave function, defined on a certain space domain $\Omega \subseteq \mathbb{R}^d$ and time interval $[t_0, T] \subset \mathbb{R}$, by

$$\Delta_\alpha = \sum_{i=1}^d \alpha_i \partial_{x_i}^2, \quad (3c)$$

$$\alpha_i \in \mathbb{R} \setminus \{0\}, \quad i \in \{1, \dots, d\}, \quad x = (x_1, \dots, x_d) \in \Omega,$$

the weighted Laplacian with respect to the spatial coordinates, by

$$V_{\beta, \gamma, \delta} : \Omega \longrightarrow \mathbb{R} : x \longmapsto \sum_{i=1}^d \left(\beta_i x_i^2 + \gamma_i \sin^2(\delta_i x_i) \right), \quad (3d)$$

$$\beta_i > 0, \quad \gamma_i, \delta_i \in \mathbb{R}, \quad i \in \{1, \dots, d\},$$

an external harmonic potential, possibly comprising an additional lattice, and by $\vartheta \in \mathbb{R}$ the coupling constant.

Underlying space domain. Quantum physics experiments concerned with the preparation and manipulation of Bose–Einstein condensates are carried out, amongst others, at the Universities of Florence and Innsbruck, see [20,30,35]. Our fundamental presumptions on the considered mathematical models, given by time-dependent Gross–Pitaevskii equations (3) or, more generally, by systems of coupled time-dependent Gross–Pitaevskii equations, see (13) below, comply with these laboratory setups. In the first instance, it is expected that the choice of the d -dimensional Euclidean space as underlying spatial domain along with asymptotic boundary conditions is well-founded, since the presence of harmonic trapping potentials ensures

$$\Omega = \mathbb{R}^d, \quad \lim_{|x| \rightarrow \infty} \Psi(x, t) = 0, \quad t \in [t_0, T]. \quad (4)$$

As detailed in Section 4, on account of a Fourier spectral space discretisation, we will replace the unbounded domain $\Omega = \mathbb{R}^d$ by a Cartesian product of bounded intervals

$$\Omega = \prod_{i=1}^d [-\omega_i, \omega_i], \quad \omega_i > 0, \quad i \in \{1, \dots, d\}, \quad (5)$$

see also (25). Notably, as solutions to (3) and (13) remain localised because of the trapping potentials, the effect of the implicitly imposed periodic boundary conditions is negligibly small as long as the intervals are chosen sufficiently large. In general, due to the fact that the confining potentials enforce exponential decay, the computation of the ground state solution allows for a valid assessment of the space intervals. More specifically, for relatively small values of the coupling constants, the first Hermite function involving an exponentially decaying factor in each spatial direction provides useful indications, see also (10) below. For relatively large values of the coupling constants, the

widths of the compactly supported Thomas–Fermi approximation are straightforwardly calculated indicators, see also (11) below.

Compact reformulation. Henceforth, we denote by $\psi : \Omega \rightarrow \mathbb{C}$ a sufficiently regular space-dependent function and employ convenient abbreviations

$$\begin{aligned} H(\psi) &= H_1(\psi) + H_2(\psi), \\ H_1(\psi) &= \Delta_\alpha \psi + V_{\beta,\gamma,\delta} \psi, \quad H_2(\psi) = \vartheta |\psi|^2 \psi, \end{aligned} \quad (6a)$$

for the operators defining the right-hand side of the partial differential equation in (3a). Consequently, we in particular obtain the compact reformulation

$$\begin{cases} \partial_t \Psi(x, t) = -i H(\Psi(x, t)), \\ \Psi(x, t_0) = \Psi_0(x), \quad (x, t) \in \Omega \times [t_0, T]. \end{cases} \quad (6b)$$

Conserved quantities. A fundamental attribute of time-dependent Gross–Pitaevskii equations is the preservation of physical quantities such as total mass (or particle number) and energy

$$\begin{aligned} M(\psi) &= \|\psi\|_{L^2}^2, \quad E(\psi) = E_1(\psi) + \frac{1}{2} E_2(\psi), \\ E_1(\psi) &= \left(H_1(\psi) \mid \psi \right)_{L^2}, \quad E_2(\psi) = \left(H_2(\psi) \mid \psi \right)_{L^2}, \end{aligned} \quad (7a)$$

see (1) and (3)–(6). Integration-by-parts under the requirement that the boundary terms vanish yields

$$E_1(\psi) = - \sum_{i=1}^d \alpha_i \left(\partial_{x_i}^2 \psi \right)^2 + \left(V_{\beta,\gamma,\delta} \mid |\psi|^2 \right)_{L^2}, \quad E_2(\psi) = \vartheta \left(|\psi|^2 \mid |\psi|^2 \right)_{L^2}, \quad (7b)$$

which reveals that the energy functional is real-valued and positive for non-trivial solutions, provided that the weights in the Laplacian are negative and that the potential as well as the coupling constant are non-negative. Straightforward calculations show that the Gâteaux derivatives are given by

$$\begin{aligned} M'(\psi) \phi &= 2 \Re(\psi \mid \phi)_{L^2} = 2 \Im(i(\psi \mid \phi)_{L^2}), \\ E'(\psi) \phi &= 2 \Re(H(\psi) \mid \phi)_{L^2} = 2 \Im(i(H(\psi) \mid \phi)_{L^2}), \end{aligned} \quad (7c)$$

see also (2). As a consequence, for regular solutions to the Gross–Pitaevskii equation (3)–(6), this implies

$$\begin{aligned} \frac{d}{dt} M(\Psi(\cdot, t)) &= M'(\Psi(\cdot, t)) \partial_t \Psi(\cdot, t) = -2 \Im \left(E_1(\Psi(\cdot, t)) + E_2(\Psi(\cdot, t)) \right) = 0, \\ \frac{d}{dt} E(\Psi(\cdot, t)) &= E'(\Psi(\cdot, t)) \partial_t \Psi(\cdot, t) = -2 \Im \left(\|H(\Psi(\cdot, t))\|_{L^2}^2 \right) = 0, \\ & \quad t \in [t_0, T], \end{aligned}$$

and thus confirms the characterisations of mass and energy conservation as

$$M(\Psi(\cdot, t)) = M(\Psi(\cdot, t_0)), \quad E(\Psi(\cdot, t)) = E(\Psi(\cdot, t_0)), \quad t \in [t_0, T]. \quad (7d)$$

Regularity and normalisation conditions. For the time-dependent Gross–Pitaevskii equation (3)–(6), due to mass conservation, it is justified to impose the normalisation condition

$$\|\Psi(\cdot, t)\|_{L^2} = \sqrt{N_0}, \quad t \in [t_0, T],$$

with a strictly positive real number $N_0 > 0$. Besides, we tacitly suppose that the prescribed initial states and hence the associated solutions are sufficiently regular such that in particular the energy functional is well-defined. Stronger regularity requirements are needed in the context of higher-order operator splitting methods to retain the full orders of convergence.

Ground and excited states. Ground and excited states are solutions to Schrödinger equations with separate dependencies on space and time variables. More precisely, inserting the ansatz

$$\begin{aligned} \Psi(x, t) &= e(t) \Phi(x), \quad (x, t) \in \Omega \times [t_0, T], \\ e : [t_0, T] &\longrightarrow \mathbb{C}, \quad |e(t)| = 1, \quad t \in [t_0, T], \\ \Phi : \Omega &\longrightarrow \mathbb{R}, \quad \|\Phi\|_{L^2} = \sqrt{N_0}, \end{aligned}$$

into the Gross–Pitaevskii equation (3)–(6), implies

$$i e'(t) \Phi(x) = e(t) H(\Phi(x)), \quad (x, t) \in \Omega \times [t_0, T],$$

and hence $e(t) = e^{-i\mu t}$ for $t \in [t_0, T]$ with $\mu \in \mathbb{R}$ as well as

$$H(\Phi(x)) = \mu \Phi(x), \quad x \in \Omega.$$

Taking the inner product and performing integration-by-parts shows the close connection of the so-called chemical potential with the energy

$$\begin{aligned} N_0 \mu &= (H(\Phi) | \Phi)_{L^2} = E(\Phi) + \frac{1}{2} E_2(\Phi) = E(\Psi(\cdot, t)) + \frac{1}{2} E_2(\Psi(\cdot, t)), \\ &t \in [t_0, T], \end{aligned}$$

and in particular confirms that it is a real-valued quantity. Summarised, the ground state solution at minimal energy level is given by a pair (Φ, μ) that satisfies the relations

$$\begin{aligned} \Psi(x, t) &= e^{-i\mu t} \Phi(x), \quad (x, t) \in \Omega \times [t_0, T], \quad \|\Phi\|_{L^2} = \sqrt{N_0}, \\ H(\Phi) &= \mu \Phi, \quad \mu = \frac{1}{N_0} \left(E(\Phi) + \frac{1}{2} E_2(\Phi) \right). \end{aligned} \quad (8)$$

Accordingly, solutions of this form at higher energy levels are referred to as excited states.

Imaginary time propagation. The imaginary time propagation, also known as normalised gradient flow method, is a heuristic approach for the computation of ground and first excited state solutions. It relies on the time integration of the parabolic counterpart of the Gross–Pitaevskii equation

$$\begin{cases} \partial_t \tilde{\Psi}(x, t) = -H(\tilde{\Psi}(x, t)), \\ \tilde{\Psi}(x, t_0) = \tilde{\Psi}_0(x), \quad (x, t) \in \Omega \times [t_0, T], \end{cases} \quad (9a)$$

formally retained from (3)–(6) by multiplying the time variable by the imaginary unit and using that $i\partial_t = -\partial_{it}$. For well-definedness, it is essential that the weights in the Laplacian are negative. In order to ensure convergence towards a non-trivial stationary solution, an additional normalisation condition is imposed, which is often realised by a straightforward scaling of the numerical solution values at certain intermediate times. For suitably chosen real-valued initial states and sufficiently large final times, it is expected that this approach yields appropriate approximations to ground and excited state solutions through

$$\sqrt{N_0} \frac{\tilde{\Psi}(x, T)}{\|\tilde{\Psi}(\cdot, T)\|_{L^2}} \approx \Phi(x), \quad x \in \Omega. \quad (9b)$$

More precisely, the resulting space-dependent function $\Phi : \Omega \rightarrow \mathbb{R}$ determines $\mu \in \mathbb{R}$ and hence $\Psi : \Omega \times [t_0, T] \rightarrow \mathbb{C}$ as stated in (8). In brief, the imaginary time propagation is substantiated by the application of the gradient descent method for the computation of the solution with minimal energy

$$\tilde{\Psi}_{\text{new}} = \tilde{\Psi}_{\text{old}} - \tau E'(\tilde{\Psi}_{\text{old}}) = \tilde{\Psi}_{\text{old}} - 2\tau H(\tilde{\Psi}_{\text{old}})$$

and its interpretation as explicit Euler approximation to (9a). Hereby, the connection between the directional derivative of the energy functional and the operator defining the right-hand side of the partial differential equation in (9a) is used, see (7c). Further enhancements of the procedure incorporate the knowledge of approximate ground and first excited state solutions in simplified settings and stepwise adaptations to the problem under consideration. Specifically, for harmonic potentials, the first Hermite function involving an exponentially decaying factor in each spatial direction

$$\begin{aligned} \gamma_i &= 0, \quad i \in \{1, \dots, d\}, \quad \vartheta = 0, \\ \Psi(x, t) &= \frac{\sqrt{N_0}}{\sqrt[4]{\pi^d}} \prod_{i=1}^d \left(\sqrt{\frac{\beta_i}{-\alpha_i}} e^{-it\sqrt{-\alpha_i\beta_i} - \frac{1}{2}\sqrt{\frac{\beta_i}{-\alpha_i}}x_i^2} \right), \\ x &= (x_1, \dots, x_d) \in \mathbb{R}^d, \end{aligned} \quad (10)$$

and the compactly supported Thomas–Fermi approximation

$$\begin{aligned} \alpha_i &= 0, \quad \gamma_i = 0, \quad i \in \{1, \dots, d\}, \\ \Phi(x) &= \begin{cases} \sqrt{W(x)}, & W(x) = \frac{1}{\vartheta} (\mu - V_{\beta, \gamma, \delta}(x)) > 0, \\ 0, & W(x) \leq 0, \end{cases} \\ \mu &= \begin{cases} \sqrt[3]{\frac{9}{16} \beta_1 \vartheta^2 N_0^2}, & d = 1, \\ \sqrt{\frac{2}{\pi} \sqrt{\beta_1 \beta_2} \vartheta N_0}, & d = 2, \\ \sqrt[5]{\frac{225}{8\pi} \beta_1 \beta_2 \beta_3 \vartheta^2 N_0^2}, & d = 3, \end{cases} \\ \Psi(x, t) &= e^{-i\mu t} \Phi(x), \quad x \in \mathbb{R}^d, \end{aligned} \quad (11)$$

yield ground state solutions for the linear Schrödinger equation and the limiting case of a negligible kinetic part, respectively. Another option to improve the convergence properties of the algorithm is the application of the gradient descent with momentum, where (9a) is replaced by

$$\begin{cases} \partial_{tt} \tilde{\Psi}(x, t) = -c \partial_t \tilde{\Psi}(x, t) - H(\tilde{\Psi}(x, t)), \\ \tilde{\Psi}(x, t_0) = \tilde{\Psi}_{10}(x), \quad \partial_t \tilde{\Psi}(x, t_0) = \tilde{\Psi}_{20}(x), \quad (x, t) \in \Omega \times [t_0, T], \end{cases} \quad (12a)$$

or an associated first-order system such as

$$\begin{cases} \partial_t \tilde{\Psi}_1(x, t) = \tilde{\Psi}_2(x, t), \\ \partial_t \tilde{\Psi}_2(x, t) = -c \tilde{\Psi}_2(x, t) - H(\tilde{\Psi}_1(x, t)), \\ \tilde{\Psi}_1(x, t_0) = \tilde{\Psi}_{10}(x), \quad \tilde{\Psi}_2(x, t_0) = \tilde{\Psi}_{20}(x), \quad (x, t) \in \Omega \times [t_0, T], \end{cases} \quad (12b)$$

with suitably chosen initial conditions for the solution and its time derivative.

3 Gross–Pitaevskii systems

Model problem. In order to exemplify systems of coupled time-dependent Gross–Pitaevskii equations describing the dynamical evolution of multi-species Bose–Einstein condensates, we consider the model problem

$$\begin{cases} i \partial_t \Psi_j(x, t) = \Delta_{\alpha_j} \Psi_j(x, t) + V_{\beta_j, \gamma_j, \delta_j}(x) \Psi_j(x, t) \\ \quad + \sum_{k=1}^J \vartheta_{jk} |\Psi_k(x, t)|^2 \Psi_j(x, t), \\ \Psi_j(x, t_0) = \Psi_{0j}(x), \quad (x, t) \in \Omega \times [t_0, T], \quad j \in \{1, \dots, J\}. \end{cases} \quad (13a)$$

By analogy to (3), the vector-valued macroscopic wave function is defined on a certain space domain and time interval

$$\Psi : \Omega \times [t_0, T] \longrightarrow \mathbb{C}^d : (x, t) \longmapsto \Psi(x, t) = \left(\Psi_1(x, t), \dots, \Psi_d(x, t) \right)^T, \quad (13b)$$

and the weighted Laplace operators and external potentials are given by

$$\begin{aligned}\Delta_{\alpha_j} &= \sum_{i=1}^d \alpha_{ji} \partial_{x_i}^2, \quad \alpha_j = (\alpha_{j1}, \dots, \alpha_{jd}), \\ V_{\beta_j, \gamma_j, \delta_j} : \Omega &\longrightarrow \mathbb{R} : x \longmapsto \sum_{i=1}^d \left(\beta_{ji} x_i^2 + \gamma_{ji} \sin^2(\delta_{ji} x_i) \right), \\ \beta_j &= (\beta_{j1}, \dots, \beta_{jd}), \quad \gamma_j = (\gamma_{j1}, \dots, \gamma_{jd}), \quad \delta_j = (\delta_{j1}, \dots, \delta_{jd}), \\ \alpha_{ji}, \beta_{ji} &> 0, \quad \gamma_{ji}, \delta_{ji} \in \mathbb{R}, \quad i \in \{1, \dots, d\}, \quad j \in \{1, \dots, J\}.\end{aligned}\tag{13c}$$

Compact reformulation. With regard to (6), we rewrite (13a) in compact form as

$$\begin{cases} i \partial_t \Psi(x, t) = H(\Psi(x, t)) \\ \Psi(x, t_0) = \Psi_0(x), \quad (x, t) \in \Omega \times [t_0, T], \end{cases}\tag{14a}$$

where we employ the convenient short notation

$$\begin{aligned}H(\psi) &= H_1(\psi) + H_2(\psi), \\ \psi &= (\psi_1, \dots, \psi_J)^T, \quad H_\ell(\psi) = \left(H_{\ell 1}(\psi), \dots, H_{\ell J}(\psi) \right)^T, \quad \ell \in \{1, 2\}, \\ H_{1j}(\psi) &= \Delta_{\alpha_j} \psi_j + V_{\beta_j, \gamma_j, \delta_j} \psi_j, \quad H_{2j}(\psi) = \sum_{k=1}^J \vartheta_{jk} |\psi_k|^2 \psi_j, \\ & j \in \{1, \dots, J\}.\end{aligned}\tag{14b}$$

Conserved quantities and ground states. By means of these abbreviations, it is straightforward to extend our considerations for a single Gross–Pitaevskii equation to systems. That is, for regular solutions to (13)–(14), total mass and energy

$$\begin{aligned}M(\psi) &= \|\psi\|_{(L^2)^d}^2, \quad E(\psi) = E_1(\psi) + \frac{1}{2} E_2(\psi), \\ E_{\ell j}(\psi) &= \left(H_{\ell j}(\psi) \mid \psi_j \right)_{L^2}, \quad E_\ell(\psi) = \left(H_\ell(\psi) \mid \psi \right)_{(L^2)^d} = \sum_{j=1}^J E_{\ell j}(\psi), \\ & \ell \in \{1, 2\}, \quad j \in \{1, \dots, J\},\end{aligned}$$

are preserved over time

$$M(\Psi(\cdot, t)) = M(\Psi(\cdot, t_0)), \quad E(\Psi(\cdot, t)) = E(\Psi(\cdot, t_0)), \quad t \in [t_0, T],$$

which justifies the normalisation condition

$$\|\Psi(\cdot, t)\|_{(L^2)^d} = \sqrt{N_0}, \quad t \in [t_0, T],$$

with a strictly positive real number $N_0 > 0$, see also (1) and (7). Ground and

first excited state solutions are characterised by the relations

$$\begin{aligned} \Psi_j(x, t) &= e^{-i\mu_j t} \Phi_j(x), \quad (x, t) \in \Omega \times [t_0, T], \quad \|\Phi_j\|_{(L^2)^d} = \sqrt{N_{0j}}, \\ H_{1j}(\Phi) + H_{2j}(\Phi) &= \mu_j \Phi_j, \quad \mu_j = \frac{1}{N_{0j}} \left(E_{1j}(\Phi_j) + E_{2j}(\Phi_j) \right), \\ N_{0j} > 0, \quad \sum_{j=1}^J N_{0j}^2 &= N_0, \quad j \in \{1, \dots, J\}, \end{aligned} \quad (15)$$

see (8). Accordingly, their numerical computation relies on the time integration of the non-reversible system

$$\begin{cases} \partial_t \tilde{\Psi}(x, t) = -H(\tilde{\Psi}(x, t)), \\ \tilde{\Psi}(x, t_0) = \tilde{\Psi}_0(x), \quad (x, t) \in \Omega \times [t_0, T], \end{cases} \quad (16a)$$

under the normalisation condition

$$\sqrt{N_0} \frac{\tilde{\Psi}(x, T)}{\|\tilde{\Psi}(\cdot, T)\|_{(L^2)^d}} \approx \Phi(x), \quad x \in \Omega, \quad (16b)$$

or a related second-order in time system, respectively, see (9) and (12).

4 Operator splitting methods

General initial value problem. In this section, to emphasise common features of our approaches for different situations, we introduce standard modified operator splitting methods for the general initial value problem

$$\begin{cases} \partial_t U(x, t) = F(U(x, t)) = F_1(U(x, t)) + F_2(U(x, t)), \\ U(x, t_0) \text{ given}, \quad (x, t) \in \Omega \times [t_0, T]. \end{cases} \quad (17)$$

Subsequently, we include detailed information concerning their realisation for Gross–Pitaevskii equations and extended systems, see also (3)–(6), (9), (12), (13)–(14), and (16).

Time-stepping approach. Due to the fact that the memory capacities needed in the time integration and imaginary time propagation of Gross–Pitaevskii systems in three space dimensions is limited, we find it most practicable to employ a time-stepping approach and to store a relatively low number of numerical solution values at once. That is, we subdivide the considered time interval in appropriately chosen subintervals and determine numerical approximation values at certain intermediate times through a recursive procedure

$$t_0 < t_1 < \dots < t_N = T, \quad U_n \approx U(\cdot, t_n), \quad n \in \{0, 1, \dots, N\}.$$

Incorporating the evident advantages of an adaptive strategy based on a local error control, we permit non-uniform temporal grid points and denote the associated time increments by

$$\tau_n = t_{n+1} - t_n, \quad n \in \{0, 1, \dots, N-1\}.$$

In view of the statement of standard and modified operator splitting methods, we express the exact solutions to (17) and the associated subproblems

$$\begin{cases} \partial_t u_j(x, t) = F_j(u_j(x, t)), \\ u_j(x, t_0) \text{ given, } (x, t) \in \Omega \times [t_0, T], \end{cases} \quad j \in \{1, 2\}, \quad (18)$$

in terms of the corresponding evolution operators

$$\begin{aligned} \mathcal{E}_{t-t_0, F}(U(\cdot, t_0)) &= U(\cdot, t), \\ \mathcal{E}_{t-t_0, F_j}(u_j(\cdot, t_0)) &= u_j(\cdot, t), \quad j \in \{1, 2\}, \\ t &\in [t_0, T], \end{aligned}$$

The analogous relations for the numerical solution values in terms of the respective splitting operators read as

$$U_{n+1} = \mathcal{S}_{\tau_n, F}(U_n) \approx U(\cdot, t_{n+1}) = \mathcal{E}_{\tau_n, F}(U(\cdot, t_n)), \quad n \in \{0, 1, \dots, N-1\}.$$

For a juxtaposition with modified operator splitting methods, we next include the general format of standard operator splitting methods and specify example methods.

Standard splitting methods. Operator splitting methods for the time integration of nonlinear evolution equations make use of the fact that the defining operators naturally decompose into two or more parts and that the numerical approximation of the associated subproblems is significantly simpler compared to the numerical approximation of the original problem, see (17) and (18). Any standard high-order splitting method is given by a composition of the form

$$\begin{aligned} U_{n+1} &= \mathcal{S}_{\tau_n, F}(U_n) \approx U(\cdot, t_{n+1}) = \mathcal{E}_{\tau_n, F}(U(\cdot, t_n)), \\ \mathcal{S}_{\tau_n, F} &= \mathcal{E}_{\tau_n, b_s F_2} \circ \mathcal{E}_{\tau_n, a_s F_1} \circ \dots \circ \mathcal{E}_{\tau_n, b_1 F_2} \circ \mathcal{E}_{\tau_n, a_1 F_1}, \\ n &\in \{0, 1, \dots, N-1\}, \end{aligned} \quad (19)$$

with suitably chosen real coefficients $(a_i, b_i)_{i=1}^s$. Prominent instances are the first-order Lie–Trotter splitting method involving a single composition per step

$$s = 1, \quad a_1 = 1, \quad b_1 = 1, \quad (20a)$$

the second-order Strang splitting method based on a symmetric composition

$$s = 2, \quad a_1 = 0, \quad a_2 = 1, \quad b_1 = \frac{1}{2} = b_2, \quad (20b)$$

and the symmetric fourth-order splitting method by YOSHIDA [37] including four compositions

$$\begin{aligned} s = 4, \quad a_1 = 0, \quad a_2 = 1 - 2b_2 = a_4, \quad a_3 = 4b_2 - 1, \\ b_1 = \frac{1}{2} - b_2 = b_4, \quad b_2 = \frac{1}{6} \left(1 - \sqrt[3]{2} - \frac{1}{2}\sqrt[3]{4} \right) = b_3. \end{aligned} \quad (20c)$$

A variety of higher-order operator splitting methods are found in the literature, see for instance [10,26]. For our numerical experiments, we select an optimised fourth-order splitting method proposed in [15] as a renowned instance with excellent properties regarding stability, accuracy, efficiency, and energy preservation over long times.

Modified operator splitting methods. As already indicated in the introduction, higher-order operator splitting methods such as (20c) necessarily involve negative coefficients. Because of severe stability issues in the context of non-reversible systems, they are thus inappropriate for ground and excited state computations. A remedy to these shortcomings of standard operator splitting methods is the application of modified operator splitting methods, which can be cast into the general form

$$\begin{aligned} U_{n+1} &= \mathcal{S}_{\tau_n, F}(U_n) \approx U(\cdot, t_{n+1}) = \mathcal{E}_{\tau_n, F}(U(\cdot, t_n)), \\ \mathcal{S}_{\tau_n, F} &= \mathcal{E}_{\tau_n, b_s F_2 + c_s \tau_n^2 G} \circ \mathcal{E}_{\tau_n, a_s F_1} \circ \cdots \circ \mathcal{E}_{\tau_n, b_1 F_2 + c_1 \tau_n^2 G} \circ \mathcal{E}_{\tau_n, a_1 F_1}, \\ n &\in \{0, 1, \dots, N-1\}, \end{aligned} \quad (21a)$$

with certain real coefficients $(a_i, b_i, c_i)_{i=1}^s$. More precisely, this kind of splitting methods relies on the efficient numerical approximation of the subproblems

$$\begin{aligned} \partial_t u_1(x, t) &= a_i F_1(u_1(x, t)), \quad u_1(x, t_n) \text{ given}, \\ \partial_t u_2(x, t) &= b_i F_2(u_2(x, t)) + c_i \tau_n^2 G(u_2(x, t)), \quad u_2(x, t_n) \text{ given}, \\ (x, t) &\in \Omega \times [t_n, t_n + \tau_n], \quad i \in \{1, \dots, s\}. \end{aligned} \quad (21b)$$

Hereby, it is remarkable that the incorporation of the operator

$$\begin{aligned} G(v) &= F_1''(v) F_2(v) F_2(v) + F_1'(v) F_2'(v) F_2(v) + F_2'(v) F_2'(v) F_1(v) \\ &\quad - F_2''(v) F_1(v) F_2(v) - 2 F_2'(v) F_1'(v) F_2(v), \end{aligned} \quad (21c)$$

where the arising derivatives are again understood as Gâteaux derivatives, see (2), permits the design of schemes such that the principal coefficients $(a_i, b_i)_{i=1}^s$ are non-negative. In conjunction with the factor τ_n^2 , this implies stability for non-reversible-in-time systems under moderate smallness assumptions on the time increments. For time-dependent Gross–Pitaevskii systems, the unitarity of the evolution operators associated with the subproblems and hence stability is ensured without additional non-negativity requirements on the real method coefficients. More detailed information on the practical realisation of modified operator splitting methods based on a Fourier spectral

space discretisation as well as their intrinsic stability properties and benefits in comparison with the alternative class of complex operator splitting methods is found at the end of this section.

Historical context and novel aspects. The alternative approach of modified operator splitting methods goes back to a seminal work of CHIN [17], where so-called modified potential or force-gradient operator splitting methods were proposed in the context of *linear* partial differential equations. In our recent contribution [12], we could extend this class of methods within a nonlinear setting such that the operator defined in (21c) results as iterated commutator $[D_{F_2}, [D_{F_2}, D_{F_1}]]$ of the associated Lie-derivatives. In the present work, we generalise the underlying idea to incorporate commutators of the defining operators to the time integration and imaginary time propagation of Gross–Pitaevskii systems. We specify (21c) for systems comprising two coupled time-dependent Gross–Pitaevskii equations (13)–(14) and their parabolic counterparts (16), which are of particular relevance in view of actual laboratory setups, see for instance [30,35].

Iterated commutators ($J = 2$). As important objectives regarding the practical implementation of modified operator splitting methods (21), we next state the iterated commutators for Gross–Pitaevskii systems involving two coupled equations. For this purpose, we use the following abbreviations for the component functions and the weighted gradients

$$\begin{aligned} J = 2, \quad \tilde{\Psi} &= (\tilde{\Psi}_1, \tilde{\Psi}_2)^T, \quad \Psi = (\Psi_1, \Psi_2)^T, \quad G = (G_1, G_2)^T, \\ \nabla &= (\partial_{x_1}, \dots, \partial_{x_d})^T, \quad \nabla_{\alpha_j} = (\alpha_{j1} \partial_{x_1}, \dots, \alpha_{jd} \partial_{x_d})^T, \quad V_j = V_{\beta_j, \gamma_j, \delta_j}, \\ & \quad j \in \{1, 2\}, \end{aligned}$$

see also (13) and (16).

- (i) We begin with the study of the parabolic-type case (16), where the first component of the operator in (21c) is given by

$$G_1(\tilde{\Psi}_1, \tilde{\Psi}_2) = 2 \left(S_0 + S_1(\tilde{\Psi}_1, \tilde{\Psi}_2) + S_2(\tilde{\Psi}_1, \tilde{\Psi}_2) + S_3(\tilde{\Psi}_1, \tilde{\Psi}_2) \right), \quad (22a)$$

$$S_0 = \nabla_{\alpha_1} V_1 \cdot \nabla V_1 \tilde{\Psi}_1, \quad (22b)$$

$$\begin{aligned}
S_1(\tilde{\Psi}_1, \tilde{\Psi}_2) &= 6 \vartheta_{11} V_1 \nabla_{\alpha_1} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_1 \tilde{\Psi}_1 \\
&\quad + 2 \vartheta_{12} V_2 \left(\nabla \tilde{\Psi}_2 \cdot \left(2 \nabla_{\alpha_1} \tilde{\Psi}_1 \tilde{\Psi}_2 + \nabla_{\alpha_1} \tilde{\Psi}_2 \tilde{\Psi}_1 \right) \right. \\
&\quad + \left. \left(\Delta_{\alpha_1} \tilde{\Psi}_2 - \Delta_{\alpha_2} \tilde{\Psi}_2 \right) \tilde{\Psi}_1 \tilde{\Psi}_2 \right) \\
&\quad + 2 \nabla_{\alpha_1} V_1 \cdot \left(3 \vartheta_{11} \nabla \tilde{\Psi}_1 \tilde{\Psi}_1 + 2 \vartheta_{12} \nabla \tilde{\Psi}_2 \tilde{\Psi}_2 \right) \tilde{\Psi}_1 \\
&\quad + 2 \vartheta_{12} \left(\nabla_{\alpha_1} V_2 \cdot \nabla \tilde{\Psi}_1 \tilde{\Psi}_2 \right. \\
&\quad + \left. 2 \left(\nabla_{\alpha_1} V_2 - \nabla_{\alpha_2} V_2 \right) \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_1 \right) \tilde{\Psi}_2 \\
&\quad - \vartheta_{11} \Delta_{\alpha_1} V_1 \tilde{\Psi}_1^3 + \vartheta_{12} \left(\Delta_{\alpha_1} V_2 - 2 \Delta_{\alpha_2} V_2 \right) \tilde{\Psi}_1 \tilde{\Psi}_2^2,
\end{aligned} \tag{22c}$$

$$\begin{aligned}
S_2(\tilde{\Psi}_1, \tilde{\Psi}_2) &= 2 \left(6 \vartheta_{11}^2 \nabla_{\alpha_1} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_1 \tilde{\Psi}_1^3 \right. \\
&\quad + 3 \left(\vartheta_{11} \vartheta_{12} + \vartheta_{12} \vartheta_{21} \right) \nabla_{\alpha_1} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_1 \tilde{\Psi}_1 \tilde{\Psi}_2^2 \\
&\quad - 2 \vartheta_{12} \vartheta_{21} \nabla_{\alpha_2} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_1 \tilde{\Psi}_1 \tilde{\Psi}_2^2 \\
&\quad + 4 \vartheta_{12} \vartheta_{22} \nabla_{\alpha_1} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_2^3 \\
&\quad + 6 \left(\vartheta_{11} \vartheta_{12} + \vartheta_{12} \vartheta_{21} \right) \nabla_{\alpha_1} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_1^2 \tilde{\Psi}_2 \\
&\quad - 4 \vartheta_{12} \vartheta_{21} \nabla_{\alpha_2} \tilde{\Psi}_1 \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_1^2 \tilde{\Psi}_2 \\
&\quad + \left(\vartheta_{12} \vartheta_{21} - \vartheta_{11} \vartheta_{12} \right) \nabla_{\alpha_1} \tilde{\Psi}_2 \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_1^3 \\
&\quad + 2 \vartheta_{12}^2 \nabla_{\alpha_1} \tilde{\Psi}_2 \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_1 \tilde{\Psi}_2^2 \\
&\quad + \left. 6 \vartheta_{12} \vartheta_{22} \left(\nabla_{\alpha_1} - \nabla_{\alpha_2} \right) \tilde{\Psi}_2 \cdot \nabla \tilde{\Psi}_2 \tilde{\Psi}_1 \tilde{\Psi}_2^2 \right),
\end{aligned} \tag{22d}$$

$$\begin{aligned}
S_3(\tilde{\Psi}_1, \tilde{\Psi}_2) &= 2 \left(2 \vartheta_{12} \vartheta_{21} \left(\Delta_{\alpha_1} - \Delta_{\alpha_2} \right) \tilde{\Psi}_1 \tilde{\Psi}_1^2 \tilde{\Psi}_2^2 \right. \\
&\quad + \left(\vartheta_{11} \vartheta_{12} - \vartheta_{12} \vartheta_{21} \right) \left(\Delta_{\alpha_2} - \Delta_{\alpha_1} \right) \tilde{\Psi}_2 \tilde{\Psi}_1^3 \tilde{\Psi}_2 \\
&\quad + \left. 2 \vartheta_{12} \vartheta_{22} \left(\Delta_{\alpha_1} - \Delta_{\alpha_2} \right) \tilde{\Psi}_2 \tilde{\Psi}_1 \tilde{\Psi}_2^3 \right).
\end{aligned} \tag{22e}$$

It is remarkable that only S_0 remains in the simplified linear case. The first and second spatial derivatives of the potentials arising in S_1 are often known explicitly. The term S_3 comprises second spatial derivatives of the solution components, whenever the weights in the Laplacians are different. For the second component, we have to exchange the index one by two and two by one, respectively.

(ii) The corresponding identity for (13) reads as

$$G_1(\Psi_1, \Psi_2) = 2 \text{i} \left(S_0 + S_1(\Psi_1, \Psi_2) + S_2(\Psi_1, \Psi_2) + S_3(\Psi_1, \Psi_2) \right) \Psi_1, \tag{23a}$$

$$S_0 = \nabla_{\alpha_1} V_1 \cdot \nabla V_1, \tag{23b}$$

$$\begin{aligned}
S_1(\Psi_1, \Psi_2) &= 2 \left(2 \vartheta_{12} \left(\nabla_{\alpha_1} V_1 - \nabla_{\alpha_2} V_2 \right) \cdot \Re \left(\nabla \Psi_2 \overline{\Psi_2} \right) \right. \\
&\quad \left. - \left(\vartheta_{11} \Delta_{\alpha_1} V_1 |\Psi_1|^2 + \vartheta_{12} \Delta_{\alpha_2} V_2 |\Psi_2|^2 \right) \right),
\end{aligned} \tag{23c}$$

$$\begin{aligned}
S_2(\Psi_1, \Psi_2) = & -2 \left(\vartheta_{11}^2 \Re(\nabla_{\alpha_1} \Psi_1 \cdot \nabla \Psi_1 \overline{\Psi_1}^2) \right. \\
& + 3 \vartheta_{11}^2 \nabla_{\alpha_1} \Psi_1 \cdot \nabla \overline{\Psi_1} |\Psi_1|^2 \\
& + 2 \vartheta_{12} \vartheta_{21} \nabla_{\alpha_2} \Psi_1 \cdot \nabla \overline{\Psi_1} |\Psi_2|^2, \\
& + 2 \vartheta_{12} \vartheta_{21} \Re(\nabla_{\alpha_2} \Psi_1 \cdot \nabla \Psi_2 \overline{\Psi_1} \overline{\Psi_2} \\
& + \nabla_{\alpha_2} \Psi_1 \cdot \nabla \overline{\Psi_2} \overline{\Psi_1} \Psi_2), \\
& - \vartheta_{12}^2 \left(\Re(\nabla_{\alpha_1} \Psi_2 \cdot \nabla \Psi_2 \overline{\Psi_2}^2) \right. \\
& + \nabla_{\alpha_1} \Psi_2 \cdot \nabla \overline{\Psi_2} |\Psi_2|^2) \\
& + 2 \vartheta_{12} \vartheta_{22} \left(\Re(\nabla_{\alpha_2} \Psi_2 \cdot \nabla \Psi_2 \overline{\Psi_2}^2) \right. \\
& + 2 \nabla_{\alpha_2} \Psi_2 \nabla \overline{\Psi_2} |\Psi_2|^2) \\
& \left. + 2 \vartheta_{11} \vartheta_{12} \nabla_{\alpha_1} \Psi_2 \nabla \overline{\Psi_2} |\Psi_1|^2 \right), \tag{23d}
\end{aligned}$$

$$\begin{aligned}
S_3(\Psi_1, \Psi_2) = & -2 \left(2 \Re(\vartheta_{11}^2 \Delta_{\alpha_1} \Psi_1 \overline{\Psi_1} |\Psi_1|^2 \right. \\
& + \vartheta_{12} \vartheta_{21} \Delta_{\alpha_2} \Psi_1 \overline{\Psi_1} |\Psi_2|^2), \\
& + 2 \Re(\vartheta_{11} \vartheta_{12} \Delta_{\alpha_1} \Psi_2 \overline{\Psi_2} |\Psi_1|^2 \\
& \left. + \vartheta_{12} \vartheta_{22} \Delta_{\alpha_2} \Psi_2 \overline{\Psi_2} |\Psi_2|^2) \right). \tag{23e}
\end{aligned}$$

The analogous identity holds for the second component, but with the index one replaced by two and two by one, respectively. Again, in the simplified linear case, only the term S_0 remains. Usually, the first and second spatial derivatives of the potentials that are present in S_1 can be calculated explicitly, whereas the first and second spatial derivatives of the solution components have to be computed numerically. However, hereby it is most remarkable that in the first component a common factor Ψ_1 and in the second component a common factor Ψ_2 can be extracted. Together with the validity of an invariance principle stated below, this special structure of the iterated commutator provides the knowledge of the exact solutions to the nonlinear subproblems on each of the temporal subintervals and hence considerably enhances the efficiency of modified operator splitting methods for time-dependent Gross–Pitaevskii systems.

Iterated commutator ($J = 1$). By setting $\alpha = (\alpha_{11}, \dots, \alpha_{1d})$, $(\alpha_{21}, \dots, \alpha_{2d}) = 0$, $\vartheta = \vartheta_{11}$, $\vartheta_{12} = \vartheta_{21} = \vartheta_{22} = 0$, $V = V_1 = V_{\beta, \gamma, \delta}$, and $V_2 = 0$ in (22) as well as (23), the corresponding identities for a single Gross–Pitaevskii equation result.

(i) On the one hand, in the context of the non-reversible equation (9), we

retain the significant simplification

$$G(\tilde{\Psi}) = 2 \left(\nabla_\alpha V \cdot \nabla V + \vartheta \left(6 V \nabla_\alpha \tilde{\Psi} \cdot \nabla \tilde{\Psi} + 6 \nabla_\alpha V \cdot \nabla \tilde{\Psi} \tilde{\Psi} - \Delta_\alpha V \tilde{\Psi}^2 \right) + 12 \vartheta^2 \nabla_\alpha \tilde{\Psi} \cdot \nabla \tilde{\Psi} \tilde{\Psi}^2 \right) \tilde{\Psi}.$$

(ii) On the other hand, in the context of the time-dependent Gross–Pitaevskii equation (3), we instead retain

$$G(\Psi) = 2i \left(\nabla_\alpha V \cdot \nabla V - 2\vartheta \Delta_{\alpha_1} V |\Psi|^2 - 2\vartheta^2 \Re(\nabla_\alpha \Psi \cdot \nabla \Psi \bar{\Psi}^2) - 6\vartheta^2 \nabla_\alpha \Psi \cdot \bar{\nabla} \Psi |\Psi|^2 - 4\vartheta^2 \Re(\Delta_\alpha \Psi \bar{\Psi}) |\Psi|^2 \right) \Psi.$$

For the special case $\alpha = (1, \dots, 1) \in \mathbb{R}^d$, both relations are consistent with equations (14) and (15) in [12].

Invariance principle. As indicated in the introduction and shortly before, an invariance principle permits the replacement of the nonlinear subproblems associated with time-dependent Gross–Pitaevskii systems by simplified linear systems with known exact solutions. In view of a compact formulation of this fundamental result, we make use of the fact that the defining operators can be rewritten as

$$H_2(\psi) = \tilde{H}_2(\psi) \psi, \quad G(\psi) = \tilde{G}(\psi) \psi.$$

Moreover, neglecting for notational simplicity the dependencies on the method coefficients $b_i, c_i \in \mathbb{R}$ and the time increment $\tau_n > 0$, we set

$$\tilde{F}_3(\psi) = -i \left(b_i \tilde{H}_2(\psi) + \tau_n^2 c_i \tilde{G}(\psi) \right), \quad i \in \{1, \dots, s\},$$

see also (21b). Our finding is that the solution to the initial value problem

$$\begin{cases} \partial_t \Psi(x, t) = \tilde{F}_3(\Psi(x, t)) \Psi(x, t), \\ \Psi(x, t_n) \text{ given, } (x, t) \in \Omega \times [t_n, t_n + \tau_n], \end{cases}$$

fulfills the simplified linear problem

$$\begin{cases} \partial_t \Psi(x, t) = \tilde{F}_3(\Psi(x, t_n)) \Psi(x, t), \\ \Psi(x, t_n) \text{ given, } (x, t) \in \Omega \times [t_n, t_n + \tau_n], \end{cases}$$

and hence is given by the explicit representation

$$\Psi(x, t) = e^{(t-t_n) \tilde{F}_3(\Psi(x, t_n))} \Psi(x, t_n), \quad (x, t) \in \Omega \times [t_n, t_n + \tau_n]. \quad (24)$$

A rigorous derivation of this relation relies on the calculation of the Gâteaux

derivative and the verification of the resulting identity

$$\begin{aligned} \frac{d}{dt} \tilde{F}_3(\Psi(\cdot, t)) &= \tilde{F}'_3(\Psi(\cdot, t)) \partial_t \Psi(\cdot, t) \\ &= \tilde{F}'_3(\Psi(\cdot, t)) \left(\tilde{F}_3(\Psi(x, t)) \Psi(x, t) \right) = 0, \quad t \in [t_n, t_n + \tau_n], \end{aligned}$$

see also (2) and (7). We here omit the technical details and instead refer to confirming numerical experiments.

Practical implementation. Modified operator splitting methods for the time integration and imaginary time propagation of Gross–Pitaevskii systems make use of the fact that both, the numerical approximation of the associated linear subproblems involving weighted Laplacians and the subproblems comprising the potentials and nonlinearities can be realised in a highly efficient manner by Fast Fourier transforms and their inverses, respectively, as well as pointwise multiplications.³

- (i) *Fourier series representations.* Employing the reasonable assumption that the prescribed initial states and hence the solutions to (13)–(14) are localised, due to the presence of harmonic potentials, and sufficiently regular, we may replace the underlying domains by Cartesian products of suitably chosen symmetric intervals

$$\Omega = \prod_{i=1}^d [-\omega_i, \omega_i], \quad \omega_i > 0, \quad i \in \{1, \dots, d\}, \quad (25)$$

see also (4) and (5). Consequently, denoting by $(\mathcal{F}_m)_{m \in \mathbb{Z}^d}$ the periodic Fourier functions

$$\begin{aligned} \mathcal{F}_m : \mathbb{R}^d \longrightarrow \mathbb{C} : x = (x_1, \dots, x_d) &\longmapsto \prod_{i=1}^d \left(\frac{1}{\sqrt{2}\omega_i} e^{i\pi m_i (x_i/\omega_i + 1)} \right), \\ m = (m_1, \dots, m_d) &\in \mathbb{Z}^d, \end{aligned}$$

and using that they form a complete orthonormal system of the Hilbert space $L^2(\Omega, \mathbb{C})$, Fourier series representations and Parseval’s identities hold true

$$\begin{aligned} \psi &= \sum_{m \in \mathbb{Z}^d} \psi_m \mathcal{F}_m, \quad \psi_m = \int_{\Omega} \psi(x) \mathcal{F}_{-m}(x) \, dx, \quad m \in \mathbb{Z}^d, \\ \|\psi\|_{L^2}^2 &= \sum_{m \in \mathbb{Z}^d} |\psi_m|^2, \quad \psi \in L^2(\Omega, \mathbb{C}). \end{aligned}$$

³ An elementary MATLAB code that illustrates the realisation of a specific modified operator splitting method for a single Gross–Pitaevskii equation is available through the publicly accessible link doi.org/10.5281/zenodo.7945624. The extension to two-component systems involves a certain amount of technical details. However, in essence, it requires the implementation of (22) and (23).

Moreover, by means of the following abbreviations for the purely imaginary eigenvalues associated with first spatial derivatives as well as the corresponding non-positive eigenvalues of second derivatives

$$\begin{aligned}
\partial_{x_j} \mathcal{F}_m &= \mu_{m_j} \mathcal{F}_m, \quad \mu_{m_j} = \frac{i\pi m_j}{\omega_j} \in i\mathbb{R}, \\
\mu_m &= \left(\mu_{m_j}\right)_{j=1}^d \in \mathbb{C}^{d \times 1}, \quad \mu_{\alpha,m} = \left(\alpha_j \mu_{m_j}\right)_{j=1}^d \in \mathbb{C}^{d \times 1}, \\
\nabla \mathcal{F}_m &= \mu_m \mathcal{F}_m, \quad \nabla_\alpha \mathcal{F}_m = \mu_{\alpha,m} \mathcal{F}_m, \\
\partial_{x_j}^2 \mathcal{F}_m &= \lambda_{m_j} \mathcal{F}_m, \quad \lambda_{m_j} = \mu_{m_j}^2 = -\frac{\pi^2 m_j^2}{\omega_j^2} \in \mathbb{R}_{\leq 0}, \\
\lambda_m &= \sum_{j=1}^d \lambda_{m_j} \in \mathbb{R}_{\leq 0}, \quad \lambda_{\alpha,m} = \sum_{j=1}^d \alpha_j \lambda_{m_j} \in \mathbb{R}_{\leq 0}, \\
\Delta \mathcal{F}_m &= \lambda_m \mathcal{F}_m, \quad \Delta_\alpha \mathcal{F}_m = \lambda_{\alpha,m} \mathcal{F}_m, \\
m &= (m_1, \dots, m_d) \in \mathbb{Z}^d, \quad x = (x_1, \dots, x_d) \in \Omega,
\end{aligned}$$

we obtain series representations such as

$$\nabla \psi = \sum_{m \in \mathbb{Z}^d} \mu_m \psi_m \mathcal{F}_m, \quad \Delta \psi = \sum_{m \in \mathbb{Z}^d} \lambda_m \psi_m \mathcal{F}_m,$$

and their analogues for weighted gradients and Laplacians.

- (ii) *Derivatives and linear subproblems.* The above identities permit to determine on the one hand the spatial derivatives of the current values of the time-discrete solutions arising in the iterated commutators and on the other hand the solutions to the linear subproblems, which can be cast into the form

$$\begin{aligned}
\frac{d}{dt} u(x, t) &= C \Delta_\alpha u(x, t), \quad u(x, t_n) \text{ given}, \\
(x, t) &\in \Omega \times [t_n, t_n + \tau_n], \quad C \in \mathbb{C}, \quad \alpha \in \mathbb{R}^d, \\
u(\cdot, t_{n+1}) &= \mathcal{E}_{\tau_n, C \Delta_\alpha} \left(\sum_{m \in \mathbb{Z}^d} u_m(\cdot, t_n) \mathcal{F}_m \right) = \sum_{m \in \mathbb{Z}^d} e^{C \tau_n \lambda_{\alpha,m}} u_m(\cdot, t_n) \mathcal{F}_m,
\end{aligned}$$

with high accuracy and efficiency by fast Fourier transforms, pointwise multiplications, and inverse fast Fourier transforms. In general, these procedures constitute the computationally most expensive components.

- (iii) *Nonlinear subproblems (Time integration).* For Schrödinger-type systems as they arise in the time integration of (13)–(14), the invariance principle stated above permits to determine the true solutions to the nonlinear subproblems by pointwise multiplications, see (24).
- (iv) *Nonlinear subproblems (Imaginary time propagation).* For parabolic-type systems (16), numerical approximations are obtained by the application of explicit Runge–Kutta methods. More concretely, in order to enhance the efficiency behaviour of fourth-order modified operator methods, we combine the second-order Strang splitting method (20b)

yielding

$$\mathcal{E}_{\frac{1}{2}\tau_n, b_i F_2} \circ \mathcal{E}_{\tau_n, c_i \tau_n^2 G_2} \circ \mathcal{E}_{\frac{1}{2}\tau_n, b_i F_2} \approx \mathcal{E}_{\tau_n, b_i F_2 + c_i \tau_n^2 G_2}, \quad i \in \{1, \dots, s\},$$

with the first-order explicit Euler method for $\mathcal{E}_{\tau_n, c_i \tau_n^2 G_2}$ and a fourth-order Runge–Kutta method $\mathcal{E}_{\frac{1}{2}\tau_n, b_i F_2}$, respectively. Whenever appropriate, for instance in the case of a single Gross–Pitaevskii equation, we may also use explicit solution representations for $\mathcal{E}_{\frac{1}{2}\tau_n, b_i F_2}$.

Example method. In our numerical experiments, we focus on a fourth-order modified operator splitting method involving a low number of compositions and a single evaluation of the iterated commutator

$$\begin{aligned} s = 3, \quad a_1 = 0, \quad a_2 = \frac{1}{2} = a_3, \quad b_1 = \frac{1}{6} = b_3, \quad b_2 = \frac{2}{3}, \\ c_1 = 0 = c_3, \quad c_2 = -\frac{1}{72}. \end{aligned} \quad (26)$$

This scheme goes back to a modified potential operator splitting method that was introduced by CHIN [17] in the context of linear problems. It can be seen as a correction of the second-order Strang splitting method (20b). Notably, the principal coefficients $(a_i, b_i)_{i=1}^s$ are non-negative and define a symmetric second-order operator splitting method.

Adaptivity in time. With regard to the evident advantages of adaptivity in time, we combine the fourth-order modified operator splitting method (26) with its basis, the second-order Strang splitting method (20b). Due to the fact that this pair of schemes is suitable for both, the time evolution and imaginary time propagation of Gross–Pitaevskii systems, this simple approach for a local error control permits to automatically adapt the time increments. As expected and indeed observed in the numerical experiments presented in Section 5, our strategy enhances stability, accuracy, and efficiency of the computations.

Stability and error analysis. As a preliminary step, it is a common assessment that the validity of nonstiff order conditions, deduced for linear ordinary differential equations, and the presumption of sufficiently regular initial states and hence solutions is adequate to retain the full order of convergence, also in substantially more complex settings of nonlinear partial differential equations. However, regarding a rigorous stability and error analysis of modified operator splitting methods of the general format (21) and the specific scheme (26), different aspects have to be taken into account.

- (i) *Mass conservation.* The conservation of mass is a fundamental property of modified operator splitting methods and related to the question of stability. In the context of time-dependent Gross–Pitaevskii systems (13), it follows from the unitarity of the evolution operators associated with the subproblems. For the linear subproblems involving weighted Laplacians,

it relies on the identity

$$\|\mathcal{E}_{\tau,i\Delta}\psi\|_{L^2} = \|\psi\|_{L^2}, \quad \tau \in \mathbb{R}, \quad \psi \in L^2(\Omega, \mathbb{C}).$$

The analogue for the subproblems comprising the remaining terms results from the above stated invariance principle (24). A crucial point is the particular structure of the iterated commutator (23), that is, the representation by purely imaginary quantities. We anticipate that mass conservation nearby machine precision has been observed in all numerical tests, wherefore this aspect is mentioned merely in a brief comment in Section 5.

- (ii) *Convergence analysis.* In our former works [32,33], we have provided a convergence analysis of standard operator splitting methods of arbitrarily high orders, see also our recent contribution on splitting methods with complex coefficients [14]. Modified operator splitting methods (21) can be seen as corrections of the underlying standard operator splitting methods, since the incorporated iterated commutators, additionally scaled by the squares of the time increments, only effect higher-order terms. Hence, it is expected that modified operator splitting methods inherit the stability properties of the underlying standard operator splitting methods, which are defined by the coefficients $(a_i, b_i)_{i=1}^s$. In conclusion, for reversible-in-time Gross–Piaevskii systems (13), no substantial stability restrictions will apply, whereas non-negative principal coefficients will be required in connection with non-reversible Gross–Piaevskii systems (16). A relevant point in the derivation of rigorous stability bounds and local error estimates is the justification that the numerical solution values remain well-defined under suitable regularity requirements on the problem data. Moreover, it is of importance to specify the optimal regularity requirements for retaining the full orders of convergence. A manuscript devoted to the numerical analysis of modified operator splitting methods for linear and nonlinear evolution equations of parabolic and Schrödinger type is currently under preparation. The provided results are also needed as prerequisites for a backward error analysis to explain the favourable preservation of energy over longer time frames.
- (iii) *Nonlinear subproblems.* For each spatial grid point, the subproblems involving the potentials, the nonlinearities, and the iterated commutators scale down to a system of coupled ordinary differential equations. In the context of reversible-in-time Gross–Piaevskii systems (13), the invariance principle (24) allows for reductions to linear Schrödinger equations with known solution representations. On the contrary, in the context of non-reversible Gross–Piaevskii systems (16), we apply explicit Runge–Kutta methods with limited stability regions for their numerical solution. Notably, the maximal values of the harmonic trapping potentials grow quadratically with the widths of the space intervals (25). In essence, stability is ensured whenever the products of the time increments and the

eigenvalues of the Jacobians of the defining right-hand sides belong to the stability regions of the respective Runge–Kutta methods. As expectable and also confirmed by our numerical tests, the use of an automatic error control is advantageous, since it ensures sufficiently small time stepsizes and thus enhances stable procedures. It is also an option to permit larger time stepsizes for the subproblems involving the weighted Laplacians and to reiterate the nonlinear subproblems.

- (iv) *Imaginary time propagation.* We recall that the imaginary time propagation or normalised gradient flow, respectively, is a heuristic approach for the numerical computation of global and local minimal energy states, the so-called ground and excited state solutions. It relies on the time evolution of non-reversible Gross–Pitaevskii systems (16) and requires an additional projection to prevent convergence of the numerical approximations towards the trivial zero solution. By its point of origin, the procedure is closely related to the gradient descent method and in turn to the first-order explicit Euler method applied to (16). In order to avoid severe instabilities and to balance stability and efficiency, common approaches use alternative time integration methods such as the first-order backward Euler method or a first-order semi-implicit Euler method, see [6]. For nonlinear systems, especially for larger values of the coupling constants, the crucial point is to ensure an energy diminishing and still advancing process. For the benefit of reliability and efficiency, we thus propose adaptive modified operator splitting methods. On the one hand, the use of an automatic error control will enhance successful ground state computations, and, on the other hand, higher-order splitting methods will provide favourable alternatives to fully implicit and semi-implicit integrators.

State-of-the-art splittings. Our recent works [11–14] are devoted to the introduction and analysis of alternative classes of operator splitting methods for linear and nonlinear evolution equations of parabolic and Schrödinger type, see also the references given therein. The design of favourable higher-order operator splitting methods is accomplished either by the consideration of complex coefficients or by the incorporation of iterated commutators. In particular, in the context of non-reversible-in-time systems, these strategies provide the possibility to overcome a second-order barrier. The first approach approves complex solutions of the order conditions, which correspond to a set of polynomial equations for the method coefficients, and offers the opportunity to find higher-order schemes with additional structural features [13]. However, due to stability restrictions and the phenomenon of order reductions for non-analytic nonlinearities as present in Gross–Pitaevskii systems (13) and hence in (16), they require deeper knowledge and attention [14]. Indeed, a higher-order complex splitting method can either fulfill the stability condition for (13) or for (16), but not both. As a consequence, it is not possible to apply the same higher-order complex splitting method for time evolution and imaginary

time propagation. For this reason, we focus on the second approach leading to so-called modified potential or force-gradient operator splitting methods [11] in the linear case and to modified operator splitting methods in the nonlinear case [12]. Specifically, the fourth-order scheme with coefficients given in (26) can be understood as a correction of the symmetric second-order Strang splitting method (20b). Its most evident characteristics are that the number of compositions is low and that the decisive coefficients are positive. This ensures that it is appropriate for the time integration of reversible-in-time as well as non-reversible systems and competitive with optimised standard splitting methods, as demonstrated in our numerical experiments, see Section 5 below. Additional numerical comparisons with complex splitting methods are found in [12]. We point out that our approach and the resulting implementation are adaptable to any other choice of the modified operator splitting method in an instant, since it suffices to redefine the method coefficients in a single component of the code.

5 Numerical experiments

Situation. In our numerical experiments, we demonstrate the favourable stability, accuracy, and long-term behaviour of adaptive modified operator splitting methods for Gross–Pitaevskii systems (13) and (16) in comparison with standard operator splitting methods. As renowned instances, we consider the first-order Lie–Trotter splitting method (20a), the second-order Strang splitting method (20b), an optimised fourth-order splitting method proposed in [15], and the fourth-order modified operator splitting method (21) for the particular choice (26). We chose the Strang splitting method, since it represents the basis of the modified operator splitting method (26) and is used for local error estimation. With regard to ground state computations, the Lie–Trotter splitting method is the simplest alternative to implicit or semi-implicit Euler methods, with significantly reduced computational effort. In the headlines of the graphics, we specify the decisive parameters, the weights in the Laplacians $C_1 = (\alpha_{ji})_{j \in \{1, \dots, J\}, i \in \{1, \dots, d\}}$, the weights in the harmonic potentials $C_2 = (\beta_{ji})_{j \in \{1, \dots, J\}, i \in \{1, \dots, d\}}$, the weights in the lattice potentials $C_3 = (\gamma_{ji})_{j \in \{1, \dots, J\}, i \in \{1, \dots, d\}}$, and the constants in the cubic nonlinearities $C_4 = (\vartheta_{jk})_{j, k \in \{1, \dots, J\}}$, see (13). We implement the time evolution from the initial time $t_0 = 0$ up to a prescribed final time T . The imaginary time propagation is stopped when either the change in energy is below the tolerance $\text{Tol} = 10^{-15}$ or the total number of iterations exceeds a certain number. We begin with numerical experiments for one-dimensional Gross–Pitaevskii equations and two-component systems involving $M = 512$ spatial grid points as well as two-dimensional Gross–Pitaevskii equations with $M = 512^2$ grid points. The most elaborate and time-consuming three-dimensional case with

$M = 100^3$ is treated in final experiments. For larger values of the coupling constants, the compactly supported Thomas–Fermi approximation (11) is an appropriate indicator for the widths of the space intervals (25). In view of our model problems, we set $\Omega = [-10, 10]^d$ by default, see Figures 11 to 13.

Stability and accuracy. In a first numerical experiment, we study reversible-in-time and non-reversible model problems reflecting the type of problems that arise in the time evolution and imaginary time propagation of Gross–Pitaevskii systems. For the purpose of verification, we make use of the fact that exact solutions are known in simplified linear cases. For the more complex nonlinear cases, numerical reference solutions are computed for refined time increments. In Figures 1 to 5, we display the global errors versus the time stepsizes in a logarithmic scale such that the slopes of the lines reflect the temporal orders of convergence. Due to the fact that the chosen initial states and hence the solutions satisfy certain regularity and consistency requirements, the nonstiff orders of convergence are indeed retained. Whenever highly accurate results are desirable, the fourth-order methods perform significantly better compared to the lower-order schemes, and the modified operator splitting method is competitive in accuracy and efficiency with the optimised standard scheme. Our main observation is that standard higher-order splitting methods applied to non-reversible systems encounter severe stability issues and even fail, whereas modified splitting methods remain stable also in higher space dimensions, provided that the time stepsizes satisfy moderate smallness assumptions. As illustrated below, these moderate stability requirements can be fulfilled by means of an inexpensive local error control based on the second-order Strang splitting method to automatically adjust the time increments.

Energy conservation. In a second numerical experiment, we illustrate the favourable behaviour of adaptive modified operator splitting methods over longer times. At first, we perform the time evolution of Gross–Pitaevskii systems by the standard splitting methods of orders two and four, respectively, for a prescribed sequence of time stepsizes. Then, we apply the fourth-order modified operator splitting method with a local error control based on the second-order Strang splitting method. We point out that the use of a local error control to automatically adapt the time stepsizes releases the users from the task to find suitable time stepsize ranges. That is, without an error control, suitable time increments had to be determined by several reruns, which involve high computational efforts, especially for the long-term evolution of Gross–Pitaevskii systems in three space dimensions. For the purpose of an initial test, we prescribe different tolerances Tol ranging from 10^{-4} to 10^{-8} and maintain a tentative approach in the sense that we leave aside the option to enlarge the time increments by additional scaling factors. Subsequently, we replace the local error estimate $\text{Err}_{\text{Local}}$ based on the difference of current approximations Ψ_{Modified} and Ψ_{Strang} by the local error estimate $\tau^2 \text{Err}_{\text{Local}}$ and repeat the

long-term evolution. For the resulting numbers of time steps N , we then once again apply the Strang splitting method and the optimised standard fourth-order splitting method with the corresponding equidistant time increments. We report that all splitting methods show an excellent mass conservation, which is a well-known intrinsic distinction to other classes of time integration methods such as Runge–Kutta and linear multistep methods. Concerning the conservation of total energy, also indicators for the achievable accuracy in the solution, the obtained results are displayed in Figures 6 to 9. We observe an excellent performance of higher-order standard and adaptive modified operator splitting methods compared to lower-order schemes. Specifically for more involved nonlinear settings, the results also suggest that adaptive modified splitting methods with the conservative strategy based on $\Psi_{\text{Modified}} - \Psi_{\text{Strang}}$ are preferable, whenever highly accurate results enclosing a favourable energy preservation are desirable, whereas efficiency is enhanced with the adapted strategy based on $\tau^2 \text{Err}_{\text{Local}}$. In order to confirm the reliability of our adaptive approach for sequences of prescribed tolerances, we include the outcomes for a Gross–Pitaevskii equation in two space dimensions and a one-dimensional two-component Gross–Pitaevskii system in Figure 10.

Imaginary time propagation and time evolution. In a final numerical experiment, we consider Gross–Pitaevskii equations in one, two, and three space dimensions for different parameter ranges. We contrast the case of a moderate constant in the nonlinearity to the case of an additional multiple lattice potential and a large constant in the nonlinearity, which is of particular relevance for physical experimental set-ups. Simplified linear cases, where the exact ground state and time-dependent solutions are known, are once again included for the purpose of validation. Moreover, it is remarkable that the knowledge of the ground state solutions and the associated chemical potentials define the time-dependent solutions $\Psi(x, t) = e^{-i\mu t} \Phi(x)$ for $(x, t) \in \Omega \times [t_0, T]$, see (15). Consequently, in modulus, the time-dependent solutions coincide with the ground states. Based on the adaptive fourth-order modified operator splitting method, we first perform the imaginary time propagation for the numerical computation of the ground state solutions and then evolve the Gross–Pitaevskii systems in time. For the linear cases, we prescribe constant initial states, and for the nonlinear cases, we determine the Thomas–Fermi approximations (11). We point out that the second-order Strang splitting method and the optimised standard fourth-order splitting method fail in the imaginary time propagation, when applied with time increment $\tau = \frac{1}{10}$, already in a single space dimension. On the contrary, the adaptive fourth-order modified operator splitting method yields reliable results in the ground state computation for the initial time stepsize $\tau = \frac{1}{10}$ and the tolerance $\text{Tol} = 10^{-5}$. The profiles of the chosen real-valued initial states for the imaginary time propagation, the resulting ground state solutions at imaginary time T , and the modulus of the time-dependent solutions evolved from $t_0 = 0$ to the final time T are illustrated in Figures 11 to 13. In space dimension three, we

determine the section along $x_3 = 0$ and display the corresponding values $\Psi(x_1, x_2, 0, t)$. As additional information and indicators for the overall accuracy, we include the total energy and the corresponding errors in the headlines of the figures. Altogether, we conclude that the adaptive fourth-order modified operator splitting method leads to reliable and accurate results in the imaginary time propagation and the time evolution of Gross–Pitaevskii equations.

6 Conclusions

Summary. In accordance with our main objective, the numerical simulation of multi-species Bose–Einstein condensates, we have devoted the present work to the introduction of adaptive modified operator splitting methods for the reliable imaginary time propagation and the efficient time evolution of two-component Gross–Pitaevskii systems. In a series of numerical tests, we have demonstrated the excellent performance of a specific adaptive fourth-order modified operator splitting method involving positive coefficients in comparison with renowned standard splitting methods. We have implemented two strategies for an inexpensive expedient local error control that are advantageous for successful computations.

Generalisations. Extensions that require minor adaptations of our approach concern more general particle interaction terms such as quintic or fractional power nonlinearities. Besides, it appears feasible to treat Gross–Pitaevskii equations with fractional Laplacians [2] by means of a straightforward calculation of the iterated commutators, neglecting possible simplifications in the resulting relations. The numerical simulation of rotating Bose–Einstein condensates under the influence of external magnetic fields, modelled by Gross–Pitaevskii equations with additional rotation terms, is based on suitable reformulations within the rotating frames. This yields non-autonomous problems of similar structures, which are resolved by the combination of Magnus-type integrators with modified operator splitting methods [4]. Another aspect of practical relevance is the imaginary time propagation of Gross–Pitaevskii systems by modified operator splitting methods combined with Hermite spectral space discretisations. As modified operator splitting methods are defined by real method coefficients and Hermite functions are real-valued, in contrast to the complex-valued Fourier functions, this approach has potential computational advantages, in particular in the context of nonlinear evolution equations with real-valued solutions.

Future investigations. Our studies in the near future will be dedicated to the efficient numerical simulation of concrete settings that arise in actual laboratory set-ups. Amongst others, this will include the investigation of strongly

interacting quantum systems driven by laser kicks and double species Bose–Einstein condensates [20,30,35]. Moreover, pursuing our theoretical findings in [14,32,33], we will provide a rigorous stability and error analysis of modified operator splitting methods for nonlinear evolution equations of parabolic and Schrödinger type.

Acknowledgements

This work has been supported by the Austrian Science Fund (FWF) through a stand-alone project (grant-doi 10.55776/PAT1281625). Fruitful discussions during a research stay at the Wolfgang Pauli Institute Vienna have inspired this work; Mechthild Thalhammer is grateful to the director Norbert Mauser and the staff members for their support and hospitality.

References

- [1] M. ANDERSON, J. ENSHER, M. MATTHEWS, C. WIEMAN, E. CORNELL. *Observation of Bose–Einstein condensation in a dilute atomic vapor*. Science 269/5221 (1995) 198–201.
- [2] X. ANTOINE, Q. TANG, J. ZHANG. *On the numerical solution and dynamical laws of nonlinear fractional Schrödinger/Gross–Pitaevskii equations*. International Journal of Computer Mathematics 95 (2018) 1423–1443.
- [3] J. AUER, E. KROTSCHECK, S. CHIN. *A fourth-order real-space algorithm for solving local Schrödinger equations*. J. Chem. Phys. 115 (2001) 6841–6846.
- [4] PH. BADER, S. BLANES, F. CASAS, M. THALHAMMER. *Efficient time integration methods for Gross–Pitaevskii equations with rotation term*. Journal of Computational Dynamics 6/2 (2019) 147–169.
- [5] W. BAO. *Ground states and dynamics of multicomponent Bose–Einstein condensates*. Multiscale Model. Simul. 2 (2004) 210–236.
- [6] W. BAO, Q. DU. *Computing the ground state solution of Bose–Einstein condensates by a normalized gradient flow*. SIAM J. Sci. Comput. 25 (2004) 1674–1697.
- [7] S. BLANES, F. CASAS. *On the necessity of negative coefficients for operator splitting schemes of order higher than two*. Appl. Numer. Math. 54 (2005) 23–37.
- [8] S. BLANES, F. CASAS. *A Concise Introduction to Geometric Numerical Integration*. CRC Press, 2016.

- [9] S. BLANES, F. CASAS, A. MURUA. *An efficient algorithm based on splitting for the time integration of the Schrödinger equation*. J. Comput. Phys. 303 (2015) 396–412.
- [10] S. BLANES, F. CASAS, A. MURUA. *Splitting methods for differential equations*. Acta Numerica 33 (2024) 1–161.
- [11] S. BLANES, F. CASAS, C. GONZÁLEZ, M. THALHAMMER. *Efficient splitting methods based on modified potentials: numerical integration of linear parabolic problems and imaginary time propagation of the Schrödinger equation*. Commun. Comput. Phys. 33 (2023) 937–961.
- [12] S. BLANES, F. CASAS, C. GONZÁLEZ, M. THALHAMMER. *Generalisation of splitting methods based on modified potentials to nonlinear evolution equations of parabolic and Schrödinger type*. Computer Physics Communications 295 (2024) 109007.
- [13] S. BLANES, F. CASAS, C. GONZÁLEZ, M. THALHAMMER. *Symmetric-conjugate splitting methods for evolution equations of parabolic type*. Journal of Computational Dynamics 11/1 (2024) 108–134.
- [14] S. BLANES, F. CASAS, C. GONZÁLEZ, M. THALHAMMER. *Splitting methods with complex coefficients for linear and nonlinear evolution equations*. Recent Progress on Numerical Analysis for Nonlinear Dispersive Equations, Chapter 3, 81–110 (2025).
- [15] S. BLANES, P. C. MOAN. *Practical symplectic partitioned Runge–Kutta and Runge–Kutta–Nyström methods*. J. Comput. Appl. Math. 142 (2002) 313–330.
- [16] M. CALIARI, S. ZUCCHER. *A fast time splitting finite difference approach to Gross–Pitaevskii equations*. Commun. Comput. Phys. 29 (2021) 1336–1364.
- [17] S. CHIN. *Symplectic integrators from composite operator factorizations*. Phys. Lett. A 226 (1997) 344–348.
- [18] S. CHIN. *Structure of positive decomposition of exponential operators*. Phys. Rev. E 71 (2005) 016703.
- [19] I. DANAILA, B. PROTAS. *Computation of ground states of the Gross–Pitaevskii functional via Riemannian optimization*. SIAM J. Sci. Comput. 39 (2017) 1102–1129.
- [20] S. DHAR, B. WANG, M. HORVATH, A. VASHISHT, Y. ZENG, M. ZVONAREV, N. GOLDMAN, Y. GUO, M. LANDINI, H.-C. NÄGERL. *Observing anyonization of bosons in a quantum gas*. Nature 642 (2025) 53–57.
- [21] A. EINSTEIN. *Quantentheorie des einatomigen idealen Gases – Zweite Abhandlung*. In: Sitzungsberichte der preussischen Akademie der Wissenschaften (1925) 3–10.
- [22] E. GROSS. *Structure of a quantized vortex in boson systems*. Nuovo Cimento 20 (1961) 454–477.

- [23] E. HAIRER, CH. LUBICH, G. WANNER. *Geometric Numerical Integration*. Springer, Second Edition, 2006.
- [24] A. ISERLES, G. QUISPTEL. *Why Geometric Numerical Integration?* In: K. Ebrahimi-Fard, M. Barbero Liñán (editors). Springer Proceedings in Mathematics & Statistics 267. Discrete Mechanics, Geometric Integration and Lie-Butcher Series, 1–28. Springer, 2018.
- [25] E. KIERI. *Stiff convergence of force-gradient operator splitting methods*. Appl. Numer. Math. 94 (2015) 33–45.
- [26] R. MCLACHLAN, G. QUISPTEL. *Splitting methods*. Acta Numerica 11 (2002) 341–434.
- [27] L. PITAEVSKII. *Vortex lines in an imperfect Bose gas*. Sov. Phys. JETP 13 (1961) 451–454.
- [28] L. LEHTOVAARA, J. TOIVANEN, J. ELORANTA. *Solution of time-independent Schrödinger equation by the imaginary time propagation method*. J. Comput. Phys. 221 (2007) 148–157.
- [29] I. OMELIAN, I. MRYGLOD, R. FOLK. *On the construction of high order force gradient algorithms for integration of motion in classical and quantum systems*. Phys. Rev. E 66 (2002) 026701.
- [30] F. RIBOLI, M. MODUGNO. *Topology of the ground state of two interacting Bose–Einstein condensates*. Phys. Rev. A 65 (2002) 063614.
- [31] J. SANZ-SERNA, M. CALVO. *Numerical Hamiltonian Problems*. Dover, 2018.
- [32] M. THALHAMMER. *High-order exponential operator splitting methods for time-dependent Schrödinger equations*. SIAM J. Numer. Anal. 46 (2008) 2022–2038.
- [33] M. THALHAMMER. *Convergence analysis of high-order time-splitting pseudo-spectral methods for nonlinear Schrödinger equations*. SIAM J. Numer. Anal. 50 (2012) 3231–3258.
- [34] M. THALHAMMER, J. ABHAU. *A numerical study of adaptive space and time discretisations for Gross–Pitaevskii equations*. J. Comput. Phys. 231 (2012) 6665–6681.
- [35] G. THALHAMMER, G. BARONTINI, L. DE SARLO, J. CATANI, F. MINARDI, M. INGUSCIO. *Double species Bose–Einstein condensate with tunable interspecies interactions*. Phys. Rev. Lett. 100 (2008) 210402.
- [36] V. VARADARAJAN. *Lie Groups, Lie Algebras, and Their Representations*. Springer, 1984.
- [37] H. YOSHIDA. *Construction of higher order symplectic integrators*. Phys. Lett. A 150 (1990) 262–268.




Fig. 1. Time evolution of linear (first column) versus nonlinear (second column) reversible-in-time model problems in one (first row) and two (second row) space dimensions. Application of standard and modified operator splitting methods, specifically the first-order Lie–Trotter splitting (20a), the second-order Strang splitting (20b), the optimised fourth-order splitting by BLANES, MOAN [15], and the fourth-order modified splitting based on (26). The slopes of the lines reflect the temporal orders of convergence and confirm the preservation of the nonstiff orders for problems with sufficiently regular solutions. Modified splitting methods remain stable and competitive in accuracy and efficiency in all test cases.




Fig. 2. Comparisons of the results displayed in Figure 1 with related two-component systems (dotted lines). The good agreement of the global errors for the cases $(J, d) \in \{(1, 1), (1, 2), (2, 1)\}$ suggests a similar behaviour regarding stability and accuracy for more complex settings in three space dimensions, see Figure 5.




Fig. 3. Time evolution of linear (first column) versus nonlinear (second column) non-reversible model problems in one (first row) and two (second row) space dimensions. Application of standard and modified operator splitting methods, see Figure 1. The slopes of the lines reflect the temporal orders of convergence and confirm the preservation of the nonstiff orders for problems with sufficiently regular solutions. For standard fourth-order splitting methods applied to non-reversible systems, severe stability issues due to the occurrence of negative method coefficients and hence even failures for larger time stepsizes are observed. On the contrary, modified splitting methods remain stable and yield highly accurate results for one space dimension (first row). For problems in two space dimensions with increased stiffness (second row), the time integration of the subproblems comprising the potentials and the nonlinearities with refined time increments (factor $\frac{1}{4}$ for Strang splitting, factor $\frac{1}{8}$ for modified splitting) leads to significant improvements.




Fig. 4. Comparisons of the results displayed in Figure 3 with related two-component systems (dotted lines). The good agreement of the global errors for the cases $(J, d) \in \{(1, 1), (1, 2), (2, 1)\}$ suggests a similar behaviour regarding stability and accuracy for more complex settings in three space dimensions, see Figure 5. Due to severe instabilities for non-reversible systems, standard higher-order operator splitting methods fail for larger time increments, whereas modified splitting methods with an appropriate resolution of the nonlinear subproblems remain stable and favourable in accuracy.




Fig. 5. Confirmation of the favourable stability and accuracy behaviour of fourth-order modified splitting methods applied to linear (left column) versus non-linear (right column) and reversible-in-time (first row) versus non-reversible (second row) model problems in three space dimensions. In order to ensure a reliable numerical approximation of the subproblems comprising the potentials and the nonlinearities for larger time stepsizes, these stepsizes are refined by a factor $\frac{1}{16}$.




Fig. 6. Energy preservation in a long-term integration of a linear Schrödinger equation. Application of standard splitting methods for prescribed numbers of time steps $N \in \{5 \cdot 10^2, 10^3, 5 \cdot 10^3, 10^4, 5 \cdot 10^4\}$ and resulting errors in the total energy (first row). Application of an adaptive modified splitting method for prescribed tolerances $\text{Tol} \in \{10^{-4}, 10^{-5}, 10^{-6}, 10^{-7}, 10^{-8}\}$ with corresponding numbers of time stepsizes and resulting errors in the total energy. On the one hand, the optimal time stepsize is determined by means of an estimate for the local error $\text{Err}_{\text{Local}}$ based on the difference $\Psi_{\text{Modified}} - \Psi_{\text{Strang}}$, which corresponds to local order three (second row, left column). On the other hand, the optimal time stepsize is determined by means of the local error estimate $\tau^2 \text{Err}_{\text{Local}}$, which corresponds to local order five and reduces the numbers of time steps (second row, right column). The quotients of subsequent tolerances $Q_1 = 10$ and the quotients of subsequent numbers of time steps, $Q_2 = 2.15$ or $Q_2 = 1.58$, respectively, indeed confirm the local orders of convergence through $\ln(Q_1)/\ln(Q_2) \approx p + 1$.




Fig. 7. Energy preservation in a long-term integration of a linear Schrödinger equation. For prescribed tolerances, the time integration is performed by an adaptive fourth-order modified operator splitting method (local error estimate $\text{Err}_{\text{Local}}$ based on $\Psi_{\text{Modified}} - \Psi_{\text{Strang}}$ for left column, local error estimate $\tau^2 \text{Err}_{\text{Local}}$ for right column). Subsequently, for the resulting numbers of time steps and corresponding equidistant time increments, standard operator splitting methods of orders two and four are applied. The results confirm the favourable performance of adaptive modified splitting methods, in particular when high accuracy is desirable.




Fig. 8. Corresponding results concerning the energy preservation in a long-term integration of a one-dimensional Gross–Pitaevskii equation, see Figure 6.




Fig. 9. Corresponding results concerning the energy preservation in a long-term evolution of a one-dimensional Gross–Pitaevskii equation, see Figure 7. For low tolerances, the conservative strategy with local error estimate $\text{Err}_{\text{Local}}$ based on $\Psi_{\text{Modified}} - \Psi_{\text{Strang}}$ (left column) leads to highly accurate results. For the adapted strategy with local error estimate $\tau^2 \text{Err}_{\text{Local}}$ and a reduced number of time steps (right column), which enhances efficiency, a connection between the errors in energy and the prescribed tolerances is observed.




Fig. 10. Corresponding results concerning the energy preservation in a long-term integration of a two-dimensional Gross–Pitaevskii equation (first row) and a one-dimensional two-component Gross–Pitaevskii system (second row), see Figure 8.




Fig. 11. Imaginary time propagation and subsequent time evolution of a one-dimensional linear Schrödinger equation (first column), a one-dimensional Gross–Pitaevskii equation with moderate nonlinearity (second column), and a one-dimensional Gross–Pitaevskii equation with additional multiple lattice potential and large constant in the nonlinearity (third column). Profiles of the prescribed initial states for the imaginary time propagation (first row). The time-dependent solutions at a short intermediate time (third row) and a larger final time (fourth row) coincide with the resulting ground state solutions at the imaginary time T (second row) in modulus. The numbers of iterations, the values of the total energies, and the corresponding errors are included in the headlines of the figures.




Fig. 12. Corresponding results for two space dimensions.




Fig. 13. Corresponding results for three space dimensions. The solution profiles along $x_3 = 0$ resemble the results obtained in the two-dimensional case, see Figure 12. Differences are in particular encountered in the number of iterations and the values of the total energies.